PHYTOPATHOLOGY

AN INTERNATIONAL JOURNAL

OFFICIAL ORGAN OF THE AMERICAN PHYTOPATHOLOGICAL SOCIETY

VOLUME 25

JANUARY–DECEMBER, 1935

EDITORS

H. B. HUMPHREY — ANNIE RATHBUN GRAVATT — EUBANKS CARSNER

EDITOR FOR EUROPE

H. M. QUANJER

ASSOCIATE EDITORS

A. G. NEWHALL	JESSIE I. WOOD	GLENN G. HAHN
ARTHUR H. EDDINS	B. F. LUTMAN	R. P. WHITE
G. B. SANFORD	LEE M. HUTCHINS	F. O. HOLMES
F. L. DRAYTON	C. D. LA RUE	J. J. CHRISTENSEN

THE AMERICAN PHYTOPATHOLOGICAL SOCIETY

OFFICERS

H. T. GÜSSOW, *President*
F. C. MEIER, *Vice-President*
H. P. BARSS, *Secretary*
H. A. EDSON, *Treasurer*

COUNCILORS

J. C. ARTHUR — F. D. HEALD — J. C. WALKER
H. B. HUMPHREY

COUNCILORS REPRESENTING DIVISIONS

T. E. RAWLINS — V. H. YOUNG

PUBLISHED FOR THE SOCIETY
THE SCIENCE PRESS PRINTING COMPANY
LANCASTER, PA.

PHYTOPATHOLOGY

AN INTERNATIONAL JOURNAL

OFFICIAL ORGAN OF THE AMERICAN PHYTOPATHOLOGICAL SOCIETY

VOLUME 25

JANUARY–DECEMBER, 1935

EDITORS

H. B. HUMPHREY — ANNIE RATHBUN GRAVATT — EUBANKS CARSNER

EDITOR FOR EUROPE

H. M. QUANJER

ASSOCIATE EDITORS

A. G. NEWHALL	JESSIE I. WOOD	GLENN G. HAHN
ARTHUR H. EDDINS	B. F. LUTMAN	R. P. WHITE
G. B. SANFORD	LEE M. HUTCHINS	F. O. HOLMES
F. L. DRAYTON	C. D. LA RUE	J. J. CHRISTENSEN

THE AMERICAN PHYTOPATHOLOGICAL SOCIETY

OFFICERS

H. T. GÜSSOW, *President*
F. C. MEIER, *Vice-President*
H. P. BARSS, *Secretary*
H. A. EDSON, *Treasurer*

COUNCILORS

J. C. ARTHUR — F. D. HEALD — J. C. WALKER
H. B. HUMPHREY

COUNCILORS REPRESENTING DIVISIONS

T. E. RAWLINS — V. H. YOUNG

PUBLISHED FOR THE SOCIETY
THE SCIENCE PRESS PRINTING COMPANY
LANCASTER, PA.

ERRATA

Page 38, WEISS abstract, line 14, *read* discomycetous *for* iscomycetous.

Page 240, heading of paragraph 3, *read* spraying *for* spring.

Page 253, line 12 *read* Tuberculina *for* Tubercularia.

Page 343, line 8, *read* from *for* for.

Page 369, line 3 from bottom of page, *read* copper *for* oil.

Page 410, line 20, last word, *read* noninoculated *for* inoculated

Page 439, line 2 *read* diseased *for* dismissed.

Page 443, line 22, *read* participle *for* principle.

Page 480, line 7, *read* virus preparation multiplied by 100, is also given *for* virus preparation, is also given.

Pages 491 and 492, heading of column 11, table 2, *read* 26.78 *for* 26.27.

Page 514, line 8, *add* All 4 forms of *T. levis* tested, infected rye.

Pages 603, line 21, *read* bandage gauze *for* cheesecloth.

Page 604, table 1, column 1, *read* $.4 \times 10^{-1}$ *for* 4×10^{-1}.

Page 856, line 3, *read* on to the surface *for* on the surface.

Page 857, line 36 *read* value *for* values.

Page 955, line 13, *read* 15 to 40 *for* 15 or 40.

Page 1115, line 15 *read* or X virus *for* of X virus.

Science
Wahr

CONTENTS OF VOLUME 25

NO. 1. JANUARY

PAGE

Abstracts of papers presented at the twenty-sixth annual meeting of The American Phytopathological Society, Pittsburgh, Pennsylvania, December 27 to 29, 1934, inclusive 1

The life history of Taphrina deformans. A. J. MIX 41

Experiments on the control of the root-knot nematode in the field with chloropicrin and other chemicals. G. H. GODFREY 67

The Raspberry White-bud disease and its relation to bitter rot of apple. KENNETH J. KADOW 91

The quality of lint and seed from cotton plants with phymatotrichum root rot. J. J. TAUBENHAUS and WALTER N. EZEKIEL 104

Bacterial leaf spot of carnations. WALTER H. BURKHOLDER and CARL E. F. GUTERMAN 114

Colletotrichum truncatum (Schw.), n. comb., on garden and Lima beans. C. F. ANDRUS and W. D. MOORE 121

Some effects of phymatotrichum root rot on microscopic characters of cotton fibers. G. M. STROMAN, J. J. TAUBENHAUS, and WALTER N. EZEKIEL 126

NO. 2. FEBRUARY

Variability of monosporic cultures of Coccomyces hiemalis. R. O. MAGIE 131

Incubation period of pea mosaic in the aphid, Macrosiphum pisi. H. T. OSBORN 160

Effect of nitrogen supply on host susceptibility to virus infection. ERNEST L. SPENCER 178

Antiseptic solutions and antiseptic adhesive tape in relation to control of hairy root, crown gall, and other overgrowths on nursery apple trees. A. J. RIKER, S. S. IVANOFF, and F. B. KILMER 192

A Stilbum disease of fig in Louisiana. EUGENE C. TIMS 208

Relation between number, size, and location of smut infections to reduction in yield of corn. I. J. JOHNSON and J. J. CHRISTENSEN 223

Dusting and spraying experiments for the control of sugar-beet leaf spot in southern Minnesota. E. L. LECLERG 234

Antagonistic action of Trichoderma on Rhizoctonia and other soil fungi. M. C. ALLEN and C. M. HAENSELER 244

Observations on Tuberculina maxima, a parasite of Cronartium ribicola. ERNEST E. HUBERT 253

Inoculations of Stagonospora curtisii on the Amaryllidaceae in California. CLAYTON O. SMITH 262

Macrosporium and colletotrichum rots of turnip roots. CHARLES CHUPP 269

Little leaf or rosette of fruit trees in California. ANTONI KOZLOWSKI 275

Phytopathological Notes

A portable rust-inoculation chamber. H. B. HUMPHREY and F. A. COFFMAN 279

Losses for corn ear rots in the United States. NEIL E. STEVENS and JESSIE I. WOOD 281

Joined spores of Ascochyta viciae. WILLARD CROSIER 283

Nuclear phenomena in Helminthosporium gramineum. T. W. GRAHAM 284

Book Review 287

NO. 3. MARCH

Dissemination of Southern celery-mosaic virus on vegetable crops in Florida. F. L. WELLMAN 289
Relation of Fusarium and Helminthosporium in barley seed to seedling blight and yield. J. J. CHRISTENSEN and E. C. STAKMAN 309
A descriptive key for plant viruses. JAMES JOHNSON and ISME A. HOGGAN 328
A leaf spot of bent grasses caused by Helminthosporium erythrospilum, n. sp. CHARLES DRECHSLER 344
The use of the quinhydrone and antimony electrodes for determining the pH of solid culture media. PAUL E. TILFORD 362
The use of oil-soluble copper as a fungicide. E. R. DE ONG 368
A bacteriophage in relation to Stewart's disease of corn. R. C. THOMAS 371
Phytopathological Notes
Rate of increase in area of apple-spur leaves. R. C. BAINES and H. M. BENEDICT 373
Hybridization between Sphaelotheca sorghi and Sorosporium reilianum. LEON J. TYLER and CLYDE P. SHUMWAY 375

NO. 4. APRIL

The host range of the Southern celery-mosaic virus. F. L. WELLMAN 377
Wojnowicia graminis as a very weak, secondary parasite of winter cereal crops. RODERICK SPRAGUE 405
Ascochyta boltshauseri on beans in Oregon. RODERICK SPRAGUE 416
The relation of soil acidity to a seedling disease in alfalfa on three Iowa soils. WALTER F. BUCHHOLTZ 421
Cytology of secondary spore formation in Ganoderma. S. R. BOSE 426
Phytopathological Notes
A probable case of sulphur starvation in tobacco. W. D. VALLEAU 430
Nonsterile soil leachate stimulating to zoosporangia production by Phytophthora sp. F. P. MEHRLICH 432
Gloeosporium leaf spot, a serious disease of orchids. CONST. J. ALEXOPOULOS 435
A new canker of hops in Oregon. G. R. HOERNER and D. C. SMITH 437
Crown gall on the Sequoia. CLAYTON O. SMITH 439
The growth of Agaricus campestris on plots treated with sodium chlorate. G. M. SHEAR 440
Air currents as a possible carrier of Ceratostomella ulmi. S. J. SMUCKER 442
Isolate vs. isolant. H. P. BARSS 443
Book Reviews 444
Constitution of The American Phytopathological Society 445
Announcement, Notes, and Notices 450

NO. 5. MAY

Aeciospore infection in Gymnoconia interstitialis by penetration of the cuticle. S. M. PADY 453
Chemical studies on the virus of tobacco mosaic III. Rates of inactivation at different hydrogen-ion concentrations. W. M. STANLEY 475
Influence of phosphorus and potassium supply on host susceptibility to yellow tobacco mosaic infection. ERNEST L. SPENCER 493
Infección experimental del centeno de Petkus (Secale cereale v. vulgare), por las caries del trigo: Tilletia tritici y Tilletia levis. RAIMUNDO NIEVES 503

Nomenclature of the tobacco downy mildew fungus. E. E. CLAYTON and JOHN A. STEVENSON 516
Phytopathological Note
Comments on the article by A. Kozlowski on little-leaf or rosette of fruit trees in California. W. H. CHANDLER and D. R. HOAGLAND 522
Book Reviews 523
Report of the twenty-sixth annual meeting of The American Phytopathological Society 528

NO. 6. JUNE

Pathologico-anatomical changes in the tomato incident to development of woodiness of the fruit. P. V. MICHAILOWA 539
The cotton root rot fungus, Phymatotrichum omnivorum, parasitic on the watermelon Citrullus vulgaris. KARL D. BUTLER 559
Fusarium yellows of celery. T. C. RYKER 578
Quantitative studies on the filtration of tobacco-mosaic virus. H. H. THORNBERRY 601
Effect of phosphate buffers on infectivity of tobacco-mosaic virus. H. H. THORNBERRY 618
The initiation of downy mildew of tobacco in North Carolina in 1934. LAWRENCE F. DIXON, RUTH A. MCLEAN, and FREDERICK A. WOLF 628
A virus of crucifers and others hosts. ISME A. HOGGAN and JAMES JOHNSON 640
Factors that influence the formation and development of mycorrhizal associations in citrus roots. H. S. REED and THERESE FREMONT 645
Phytopathological Notes
Breeding millet resistant to smut in North China. CHIH TU and H. W. LI 648
An example of spread of veinbanding from potatoes to tobacco. E. M. JOHNSON 650
A root rot of Chinese elms. HOWARD LAMB, ERNEST WRIGHT, and ROSS W. DAVIDSON 652
Mites on Potatoes. DONALD REDDICK 654
Book Review 655

NO. 7. JULY

Comparative studies of certain cultures of Puccinia rubigo-vera and Puccinia tomipara on wild grasses. GEORGE W. FISCHER 657
Serological evidence in plant-virus classification. K. STARR CHESTER 686
The antigenicity of the plant viruses. K. STARR CHESTER 702
The effect of light on the initiation of rust infection. HELEN HART and I. L. FORBES 715
Phytopathological Notes
An internal necrosis of bean seeds. C. R. ORTON and W. D. HENRY 726
Filtrability of certain plant pathogenic bacteria. P. A. ARK 728
Soggy breakdown of winter banana apples. H. H. PLAGGE and T. J. MANEY 730

NO. 8. AUGUST

Fusarium strains in relation to wilt of China aster. REGINA S. RIKER and L. R. JONES 733
Rhizoctonosis of potatoes grown under irrigation. L. A. SCHAAL 748
The nature of inhibition between certain fungi parasitic on citrus. JAIME GUISCAFRÉ ARRILLAGA 763
Acquired immunity from cucumber mosaic in zinnia. W. C. PRICE 776
Morphologic studies of Septoria lycopersici. HUBERT A. HARRIS 790

A phytophthora disease of peach seedlings. JOHN C. DUNEGAN 800
Summary of investigations with Ustilago striaeformis parasitizing some common grasses. W. H. DAVIS 810

NO. 9. SEPTEMBER

Resistance of Pinus sylvestris to a gall-forming Peridermium. W. G. HUTCHINSON 819
Graft versus insect transmissions of curly top in tomatoes (tomato yellows). MICHAEL SHAPOVALOV 844
The influence of Phytomonas tumefaciens and Phytomonas rhizogenes on the actual acidity of certain liquid and agar substrata. A. R. WILSON 854
Effect of certain chemicals on the "combination streak" virus of tomatoes. MICHAEL SHAPOVALOV 864
The inheritance of resistance to common bean mosaic in field and garden beans. W. H. PIERCE 875
Phytopathological Notes
Areolate spot of citrus caused by Leptosphaeria bondari. A. A. BITANCOURT and ANNA E. JENKINS 884
Transfer of Diaporthe umbrina to the genus Cryptosporella. ANNA E. JENKINS and LEWIS WEHMEYER 886
Report of the Nineteenth Annual Meeting of the Pacific Division of the American Phytopathological Society 890

NO. 10. OCTOBER

Chemical studies on the virus of tobacco mosaic. IV. Some effects of different chemical agents on infectivity. W. M. STANLEY 899
Chemical studies on the virus of tobacco mosaic. V. Determination of optimum hydrogen-ion concentrations for purification by precipitation with lead acetate. W. M. STANLEY 922
Effect of tannic acid on the infectivity of tobacco-mosaic virus. H. H. THORNBERRY 931
Particle diameter of certain plant viruses and Phytomonas pruni bacteriophage. H. H. THORNBERRY 938
Classification of Southern celery-mosaic virus. W. C. PRICE 947
Phytopathological Notes
Peronospora in storage cabbage. G. B. RAMSEY 955
Toxicity of low concentrations of ammonia to mycelium and sclerotia of Sclerotium rolfsii. L. D. LEACH and A. E. DAVEY 957
Blackberries: Possible source of streak infection in black raspberries. J. K. THORNTON 959
Susceptibility of raspberry species and varieties to leaf spot (Mycosphaerella rubi) at Beltsville, Maryland. GEORGE M. DARROW 961
Sclerotinia wilt of Canada thistle. E. W. BODINE 963
Report of the 1935 annual meeting of the Southern Division of The American Phytopathological Society 965

NO. 11. NOVEMBER

Rust resistance in Antirrhinum. E. B. MAINS 977
Studies on the host range of Phytomonas stewarti and P. vascularum. S. S. IVANOFF 992
New economic hosts of the stem-and bulb-infesting nematode. G. H. GODFREY and C. EMLEN SCOTT 1003

Cultural variations of Thielaviopsis basicola. E. M. JOHNSON and W. D. VALLEAU 1011
Factors affecting experimental error in greenhouse pot tests with sugar beets. E. L. LECLERG 1019
The demonstration of plant parasitic nematodes in host tissues. J. CEDRIC CARTER 1026
Cultural studies on the virus of tobacco mosaic. FRANK H. JOHNSON 1035
Phytopathological Notes
An effective and easily applied method of inoculating seed barley with covered smut. V. F. TAPKE 1038
The transmission of Ceratostomella ulmi through root grafts. A. F. VERRALL and T. W. GRAHAM 1039

NO. 12. DECEMBER

The effect of light intensity and temperature on infection of Hope wheat by Puccinia graminis tritici. HELEN HART and KAROL ZALESKI 1041
Studies on frenching of tobacco. ERNEST L. SPENCER 1067
A study of coalescing haploid pustules in Puccinia helianthi. A. M. BROWN 1085
Studies on seven differentiating characteristics of two physiologic forms of Tilletia tritici. C. S. HOLTON 1091
Effect of crown gall, hairy root, and woolly aphids on apple trees in the orchard. C. D. SHERBAKOFF and J. A. MCCLINTOCK 1099
Dates of production of the different spore stages of Cronartium ribicola in the Pacific Northwest. J. J. MIELKE and J. W. KIMMEY 1104
Seed transmission of Helminthosporium of corn. W. D. VALLEAU 1109
Phytopathological Notes
Some factors affecting infection with and spread of Diplodia zeae in the host tissue. J. R. HOLBERT, P. E. HOPPE, and A. L. SMITH 1113
A top-necrosis virus found in some apparently "healthy" potatoes. T. P. DYKSTRA 1115
An apple canker caused by Monochaetia mali. JOHN W. ROBERTS 1116
Notes on the occurrence of Tuberculina maxima on the aecia of Cronartium cerebrum. GEORGE G. HEDGCOCK 1117

INDEX TO VOLUME 25

New species in blackface type

Acetate, lead, determination of optimum hydrogen-ion concentrations with, 922–930

Acidity, of certain liquid and agar substrata, influence of Phytomonas tumefaciens and P. rhizogenes on, 854–863

Actinomyces scabies, nature of resistance to, 13; reaction of potato seedlings of inbred lines to, 13–14

Adenostegia pilosa, highly resistant to snapdragon rust, 985

Aerial Rhizoctonia on beans, 38

Aeroplane, dusting with sulphur to combat stem rust of wheat, 892–893; exposure of microscopic slides from, 27

Agaricus campestris, growth of, on plots treated with sodium chlorate, 440–442

Agent, inhibitory, produced by fungi, diffusible nature of, 1031–1034

Agropyron, inerme, Wojnowicia graminis on, 408; riparium, W. graminis on, 408; spp., inoculation experiments with Puccinia rubigo-vera, 663–671

Agrostis, alba, leaf spot of, caused by **Helminthosporium erythrospilum**, n. sp., 344–361; palustris, leaf spot of, caused by **H. erythrospilum**, n. sp., 345–361; palustris, physiologic form of Ustilago striaeformis on 816; tenuis, leaf spot of, caused by **H. erythrospilum**, n. sp., 345–361

Air, currents, as possible carrier of Ceratostomella ulmi, 442–443; microorganisms in, 27

Albino strain, of barley smut, 27–28; of Ustilago hordei, 27–28

ALEXOPOULOS, CONST. J., Gloeosporium leaf spot, a serious disease of orchids, 435–437

Alfalfa, damping off, caused by pythiaceous fungi, 8; damping off, on acid and neutral soils, 8–9; damping off, treatment of soil, 9; seedling disease of, related to soil acidity, 421–425

ALLEN, M. C., and C. M. HAENSELER, Antagonistic action of Trichoderma on Rhizoctonia and other soil fungi, 244–252

ALLISON, C. C., Hybridization between Ustilago hordei and U. medians, 5; see MOORE, M. B.

Allium, cepa, susceptible to celery virus 1, 381, 384, 394; sativum, nematode of, longevity in soils, 1009, 1010

Alternaria, associated with bottom rot of calla rhizomes, 39; in barley seed, commonest cause of seed blight, 310; sp., in corn seed, 1111, 1112

Aluminum effect of, on prevention of and recovery from tobacco frenching, 1075–1078

Amaryllidaceae, inoculations of Stagonospora curtisii on, 262–268; list of, inoculated with S. curtisii, 266, 267

Amaryllis belladonna, infection with Stagonospora curtisii, 264; belladonna, S. curtisii occasional on, 263; host for spotted wilt in California, 17

Ambrosia elatior, host of Southern celery-mosaic virus, 289–308; spread of Southern celery-mosaic virus on, in Florida, 299–308; susceptible to celery virus 1, 389, 390, 398

American Phytopathological Society, Constitution, 445; report of, annual meeting of Southern Division, 965; report of, nineteenth annual meeting of Pacific Division, 890–891; report of, Summer meeting, 451–452; report of, twenty-sixth annual meeting, 528–538

Ammonia, nitrogen, effect of, on growth of cotton-root-rot fungus, 967–968; toxicity of compounds of, to Sclerotium rolfsii, 895–896; toxicity of low concentrations of, to mycelium and sclerotia of S. rolfsii, 957–959

Anaerobic soil conditions cause of little leaf or rosette of fruit trees in California, 275–278

Ananas sativus, cripple and dark stripe, in Hawaii, 25–26; effect of Pseudococcus brevipes, 10; see pineapple; virus, 10

ANDERSON, H. W., Black stele root rot of strawberry, 5; see KADOW, KENNETH J.

ANDES, J. O., Black root rot of apple, 976; Tobacco bed soil disinfection, 972
ANDRUS, C. F., and W. D. MOORE, **Colletotrichum truncatum**, n. comb., on garden and Lima beans, 121–125
Anemone virginiana inoculation experiments with Puccinia rubigo-vera, 667
Anethum graveolens susceptible to celery virus 1, 381, 395
Angiopomopsis lophostoma, comparison of, with Wojnowicia graminis, 412
Anguillulina dipsaci, demonstration of, in garlic plants, 1027, 1028; experimental inoculations with, 1007–1009; new economic hosts of, 1003–1010
Angular leaf spot of cotton in Mississippi, 973
Antagonism of Trichoderma on Rhizoctonia and other soil fungi, 244–252
Anthracnose, of eggplants, 967; of strawberry, caused by Colletotrichum fragariae, 973–974; on Ribes, 6
Antibodies as modified antigens, 697
Antigenicity, of the plant viruses, 702–714; of viruses, 702–714
Antimony electrode, use of 362–367
Antirrhinum, glutinosum, used in crosses for rust resistance, 985; majus, infected by rust, 977; majus, infection with Puccinia antirrhini, effect of light on, 715–725; majus, susceptible to celery virus 1, 386, 388, 398; pollination technic, 980; resistance of, to rust, 977–991; rust, resistance to, 977–991; species, reaction to rust, 986; species resistant to snapdragon rust, 985; species, susceptible to snapdragon rust, 985
Antiseptics, control of hairy root and crown gall, 192–207; hairy root, crown gall, and crown gall control, list of, 195
Aphanomyces spp. on roots of spinach and flax, 14–15
Aphia rumicis, carrier of pea mosaic, tested, 174
Aphid, incubation of virus, of pea mosaic in, 31, 160–177; potato, Macrosiphum gei as carrier of pea mosaic, 174; transmission of virus of pea mosaic by, 31
Aphids, as vectors of Southern celery-mosaic virus among perennial and annual hosts, 307; transmission of cabbage mosaic by, 6; woolly, effect or, on apple trees, 975; woolly, effect of, on apple trees in orchard, 1099–1103
Aphis gossypii, transmission of Southern celery-mosaic virus by, 951; vector of celery-mosaic viruses, 891; vector of Southern celery-mosaic virus among several wild and cultivated hosts, 290–308
Apium graveolens, infected with Anguillulina dipsaci, 1006; see celery; spread of Southern celery-mosaic virus on, 290–308
Aplanobacter stewarti, bacteriophage in relation to 371–372; insect vector of, 32; see Phytomonas
Apothecium production in Sclerotinia trifoliorum and S. sclerotiorum, 19–20
Apple, bitter rot, relation to raspberry white-bud disease, 93–100; black root rot of, 976; canker caused by Monochaetia mali, 1116–1117; effect of fungicide on growth of, as factor in comparative tests, 15–16; Erwinia amylovora, relation of carbohydrate-nitrogen nutrition to infection by, 26; hairy root, environmental factors influencing development of, 32; mites on, 654; Physolospora cydoniae, variation in isolates, 16–17; relation of host vigor to infection by Xylaria mali, 12–13; rust, nature of resistance to, 30; rust, resistance of varieties, cytological study of, 30; scab, control by copper phosphate mixture, 32; scab, leaf growth rate in relation to control of, 373–375; soggy breakdown of, 730–731; spur leaves of, rate of increase in area of, 373–375; trees, effect of crown gall on, 975; trees, effect of hairy root on, 975; trees, effect of woolly aphids on, 975; trees in orchard, effect of crown gall on, 1099–1103; trees in orchard, effect of hairy root on, 1099–1103; trees in orchard, effect of woolly aphids on, 1099–1103; trunk canker, caused by Phytophthora cactorum, 5; trunk canker, control, 5; trunk canker, destructive disease in Indiana, 5; trunk canker, symptoms of, 5; trunk canker, varietal susceptibility, 5; Xylaria mali, 12

Apples, bitter rot of, relation to raspberry white-bud disease, 91–112

Areolate spot of citrus caused by **Leptosphaeria bondari, n. sp.**, 884–886

ARK, P. A., Filtrability of certain plant pathogenic bacteria, 728–729

ARMSTRONG, G. M., and C. N. CLAYTON, The reaction of several isolations of the cotton-wilt fungus to toxic dyes, 970–971; and C. B. Sumner, Studies on downy mildew of tobacco, 971

ARNDT, C. H., The etiology of damping off of cotton seedlings, 968–969; A resumé of cotton seed treatments in South Carolina, 970

ARRILLAGA, JAIME GUISCAFRÉ, Nature of inhibition between certain fungi parasitic on citrus, 763–775

ARTSCHWAGER, ERNST, book review, Handbuch der Pflanzenkrankheiten, Vol. I. Die Nicht parasitären und Virus-Krankheiten. Part II. (Nonparasitic and virus diseases) by Paul Sorauer, 525–527

Ascochyta, boltshauseri on beans in Oregon, 416–420; viciae, fused pycnospores of, 283–284

Ascospora pisi, synonym of Ascochyta bolthauseri, 419

Asexual spores of Pleurotus corticatus, 23

Aspergillus, flavus, on dent corn, 24; niger, inhibitory effect on other fungi, 9; tamarii, on dent corn, 24

Aster, China, Fusarium strains in relation to wilt of, 733–747; China, root rot of, in California, caused by Phytophthora cryptogea, 895; leaf rust, control, 17; leaf rust, varietal susceptibility to, 17–18; yellows, retention by aphids, 175

ATANASOFF, D., Bolesti na kulturnitĕ rastenija. (Diseases of cultivated plants), book review, 444; Virus diseases of plants. A bibliography, book review, 444

Atropa belladonna susceptible to celery virus 1, 386, 397

Azalea indica fungus spot of flowers in South Carolina, 38

Bacillus, aerogenes, effect of, on tobacco-mosaic virus, 33; proteus, effect of, on tobacco-mosaic virus, 33; symbiont of Pseudococcus brevipes, relation to pineapple green spotting, 10

Bacteria, Aplanobacter stewarti, bacteriophage in relation to, 371–372; associated with discoloration in Platanus sp., 19; associated with hop canker, 437–439; associated with vascular disease of poplar and willow, 18–19; cause of sweet pea fasciation, 36; effect of, mixed cultures on tobacco-mosaic virus, 33; inhibitory effect on fungi, 9; pathogenic, inducing cell stimulation in plants, physiologic studies of, 31; soft-rot and colon-typhoid-dysentery groups, pathogenicity of, 19; soft-rot and colon-typhoid-dysentery groups, physiology of, 34–35

Bacterial, blight of barley seed, 310, 325; canker of tomato, control of, 968; leaf blight of corn, effect of, on invasion by Diplodia zeae, 1113; leaf spot, of carnations, 114–120; leaf spot of cotton, dissemination of organism, 971; parasite of stem rust, 1053; spot of tomato, control of, 968; wilt of corn and its insect vectors of, 32; wilt of corn on teosinte, 32; wilt of corn, resistant lines and hybrid, 21–22; wilt of potatoes, soil treatment with sulphur and limestone, 16

Bacteriophage, in relation to Stewart's disease of corn, 371–372; Phytomonas pruni, particle diameter of certain plant viruses and, 938–946; relation of, to Stewart's disease of corn, 371–372

Bacterium, angulatum, epidemiology, 11; cactivorum, on cactus in Italy, 9; malvacearum, dissemination of, 971; solanacearum, on potato in Florida, control of, 16; tabacum, epidemiology, 11; tabacum, on Physalis virginiana, 6; translucens undulosum, effect of, on stem-rust infection, 1053

BAINES, R. C., Phytophthora trunk canker of apple, 5; and H. M. BENEDICT, Rate of increase in area of apple-spur leaves, 373–375

BANFIELD, W. M., and E. C. MANDENBERG, Importance of sanitation in controlling crown gall of the red raspberry, 5–6

Barberry eradication in relation to decreasing severity of stem rust, 25

Bark, beetles, isolation of Ceratostomella ulmi from, 37–38; disease of Tahiti lime trees caused by Phomopsis citri and Diplodia natalensis, 966

Barley, scab, caused by Gibberella saubinetii, recent outbreaks of, 309; seed, bacterial blight of, 310, 325; seed, blight of, 309–326; seed, blight of, caused by Alternaria, 310; seed, cultures from, 310; seed, disinfection of, 310; seed, fungous blight of, 310, 325; seed, microflora of, 310–313; seed, relation of Fusarium in, to discoloration of coleoptile, 313–318; seed, relation of Fusarium in, to germination, 313–318; seed, relation of Fusarium in, to root rot, 313–318; seed, relation of Fusarium in, to seedling blight, 309–327; seed, relation of Fusarium in, to stand, 313–318; seed, relation of Fusarium in, to stunting, 313–318; seed, relation of Fusarium in, to yield, 309–327; seed, relation of Helminthosporium in, to discoloration of coleoptile, 313–318; seed, relation of Helminthosporium in, to germination, 313–318; seed, relation of Helminthosporium in, to root rot, 313–318; seed, relation of Helminthosporium in, to seedling blight, 309–327; seed, relation of Helminthosporium in, to stand, 313–318; seed, relation of Helminthosporium in, to stunting, 313–318; seed, relation of Helminthosporium in, to yield, 309–327; seed, treatment of, for Fusarium, 318–326; seed, treatment of, for Helminthosporium, 318–326; seedling blight of, effect of seed treatment on, 318–326; smut, albino strain, 27–28; smut, intermediate types, distribution of, 28; susceptibility to hybrids between Ustilago hordei and U. medians, 5; susceptibility, to U. hordei, 5; susceptibility, to U. medians, 5; U. medians on, distribution of, 28; Wojnowicia graminis on, 408

Barrett, J. T., An explanation of Rhiomyxa hypogeae, 898; and Horace Parker, Observations on a species of Ligniera, a root parasite of Stellaria media, 893

Barss, H. P., Isolate *vs.* isolant

Bartholomew, E. T., Endoxerosis of lemon fruits as affected by the application of different amounts of irrigation water, 897

Beach, W. S., Tobacco wildfire, control, in Pennsylvania, 6

Bean, Ascochyta bolthauseri on, in Oregon, 416–420; **Colletotrichum truncatum**, n. comb., on, 121–125; Lima, mosaic, control with sulphur, 36; Lima, Rhizoctonia sp. on, in Florida, 38; mites on, 654; mosaic, control with sulphur, 36; mosaic, effect on transpiration, 18; Rhizoctonosis sp. in on, in Florida, 38; rust, host-parasite relationship, 39; var. Early Golden Cluster, as test plant, tobacco virus inactivation, 477

Beans, effect of tannic acid on the infectivity of tobacco-mosaic virus, 931–937; field and garden, inheritance of resistance to common bean mosaic, 875–883

Bechenbach, J. R., see Young, H. C.

Beet, leaf hopper, experiments on the multiplication of curly-top virus in, 891; sugar, effect of carbon dioxide on the pH gradient of, 891; sugar, factors affecting experimental error in greenhouse pot tests, 1019–1025; sugar, Sclerotium rot of, 895; susceptible to Southern celery mosaic, 381, 384

Beets, retention of mosaic virus by aphids, 175; sugar, soil amendments for Southern sclerotium rot of, 896

Begonia host for spotted wilt in California, 17

Benedict, H. M., see Baines, R. C.

Beta vulgaris, cicla, susceptible to celery virus 1, 381, 395; damping off, influence of, preceding crops on, 13; susceptible to celery virus 1, 381, 384, 395

Biochemic studies, on composition of crown gall, 11; on the metabolism of crown-gall and hairy-root bacteria, 11

Bitancourt, A. A., and Anna E. Jenkins, Areolate spot of citrus caused by **Leptosphaeria bondari**, n. sp., 884–886

Bitter rot of apples, relation to raspberry white-bud disease, 91–112
Black, -locust seedlings, phytophthora wilt, control, 25; locust, seedlings, phytophthora wilt, in Virginia, 24; -locust seedlings, phytophthora wilt, symptoms, 24–25; root rot, of apple, 976; stele root rot, on strawberry, 5
Blackberries, possible source of streak infection in black raspberries, 959–961
Blackberry, Gymnoconia interstitialis on aeciospore infection by, 454–466
BLANK, L. M., A mosaic on cabbage in Wisconsin, 6
Blight, new, of pea in California, 891–892; of barley seed, 309–326; Oregon tip, of tomato, interpretation on basis of viruses causing, 897–898; seedling, of barley, effect of seed treatment on, 318–326; seedling, of barley, relation of Fusarium in seed to, 309–327; seedling, of barley, relation of Helminthosporium in seed to, 308–327; stem, of Lespedeza sericea, 976
BLISS, DONALD E., The relation of Penicillium vermoeseni to a disease of ornamental palms, 896
BLODGETT, E. C., Pathogenicity and physiology of Pseudopeziza ribis, 6–7
BODINE, E. W., Sclerotina wilt of Canada thistle, 963–964
Book reviews, Bolesti na kulturnitě rastenija (Diseases of cultivated plants), 444; Handbuch der Pflanzenkrankheiten, Volume I, Part II, sixth edition. Die Nichtparasitären und Virus-Krankheiten (Non-parasitic and virus diseases), 525–527; Manual of diseases of new bast fiber plants, (Russian), 655; Plant parasitic nematodes, 287; Root rots and degeneration of the strawberry, 523–535; Virus diseases of plants: A bibliography, 444
Bordeaux, mixture, absorption by conidia of Sclerotinia fructicola, 17; mixture direct effect of, on early cucumber production, 974; mixture, for control of Macrosporium blight of tomato, 968; mixture, insoluble copper compounds as substitutes for, 40; mixture, used in sugar-beet leaf-spot control, 234–243; mode of action of, on Mycosphaerella fragariae, 970; use of, as fungicide, compared with oil-soluble copper spray, 368–370
BOSE, S. R., Cytology of secondary spore formation in Ganoderma, 426–429
Bottom rot of calla rhizomes, fungi associated with, 39
BOYCHENKO, E., see GITMAN, L.
BOYD, O. C., Evidence of the seed-borne nature of late blight (Phytophthora infestans), of tomatoes, 7
BRAMBLE, W. C., and E. C. HOLST, Microorganisms infecting pines attacked by Dendroctonus frontalis, 7
Brassica, oleracea, var. capitata, mosaic, 6; rapa, susceptible to celery virus, 1, 381, 395
Breeding peanuts for disease resistance, 971–972
Brevicoryne brassicae, vector of virus of crucifers, 640
BRIERLEY, PHILIP, Streak, a virus disease of roses, 7–8; Symptoms of rose mosaic, 8
Bromus, spp., inoculation experiments, with Puccinia rubigo-vera, 663–671; spp., inoculation experiments with P. tomipara, 674–679; tectorum, Wojnowicia graminis on, 408
BROOKS, A. N., A rhizoctonia bud rot of strawberry, 965–966; Anthracnose and wilt of strawberry caused by Colletotrichum fragariae, 973–974
Browallia, host for spotted wilt in California, 17
BROWN, A. M., A study of coalescing haploid pustules in Puccinia helianthi, 1085–1090
BRUYN, HELENA L. G. de, Heterothallism in Peronospora parasitica, 8
BUCHHOLTZ, W. F., Damping off of alfalfa on acid and neutral Iowa soils, 8-9; Relation of soil acidity to a seedling disease of alfalfa on three Iowa soils, 421–425
Bud rot, Rhizoctonia, of strawberry, 965–966

Buffers, effect on virus of tobacco mosaic, 912; phosphate, effect on infectivity of tobacco-mosaic virus, 618–627
Bulb-infesting nematode, new hosts of, 1003–1010
Bunt, wheat, experimental infection, of rye, 503–515
BURKHOLDER, WALTER H., and CARL E. F. GUTERMAN, Bacterial leaf spot of carnations, 114–120
BUTLER, KARL D., Cotton root rot fungus, Phymatotrichum omnivorum, parasitic on watermelon, Citrullus vulgaris, 559–577
BUZZATI-TRAVERSO, A., Six microorganisms pathogenic on cactus, 9

Cabbage, mosaic, effect on seed production, 6; mosaic, in Wisconsin, 6; mosaic, symptoms, 6; mosaic, transmission of, 6; Peronospora parasitica on, in storage, 955–957
Cactus, maxonii, disease of, 9; microorganisms pathogenic on, 9
Cadmium chloride, antiseptic, tests in control of hairy root and crown gall, 194–207
California celery mosaic diseases, 891
Calla bottom rot of rhizomes, 38–39; fungi associated with, 39
CALLENBACH, J. A., see COOK, H. T.
Callistephus, chinensis, Fusarium strains causing wilt of, 733–747
Campanula, host for spotted wilt in California, 17
Canker of hops in Oregon, 437–439
Capsicum, annuum, Southern celery-mosaic virus on, 304–305; annuum, susceptible to celery virus 1, 397; frutescens, susceptible to celery virus 1, 381, 382, 397
Carbohydrates, serically active, of genotypes of Neurospora tetrasperma and N. sitophila, 39
Carbon, bisulphide, control of root knot with, 76–86; compounds, effect on variability of Coccomyces hiemalis, 147, 148; dioxide, effect on the pH gradient of the sugar beet and on the feeding of the curly-top vector, 891
Carnation bacterial leaf spot, caused by Phytomonas woodsii, 118–119; control, 119; symptoms, 115
Carotenoid pigments of urediospores of Puccinia graminis, 30
Carrot susceptible to Southern celery mosaic, 381
CARTER, J. CEDRIC, Diffusible nature of the inhibitory agent produced by fungi, 1031–1034; Growth association of microorganisms, 9
CARTER, WALTER, Mechanical transmission of two viruses to pineapple, 10; The symbionts of Pseudococcus brevipes in relation to a phytotoxis secretion of the insect, 10
CASSELL, ROBERT C., see STAKMAN, E. C.
Cauliflower, host for spotted wilt in California, 17; Phytophthora megasperma causing root rot of, in California, 893–894
Celeriac susceptible to Southern celery virus, 381
Celery, calico, in California, 891; fusarium wilt, forms of, 29; fusarium yellows, 578–600; host for spotted wilt in California, 17; infected with Anguillulina dipsaci, 1006; mosaic diseases of, in California, 891; mosaic, dissemination of, on vegetable crops in Florida, 289–308; mosaic, spread of, in seed beds, 302; mosaic, Western, in California, 891; relation of soil temperature and moisture to yellows development, 585–590; Southern mosaic virus, classification of, 947–954; Southern mosaic virus, host range of, 377–404; varietal resistance to yellows, 590–593; virus 1, cause of Southern celery mosaic, dissemination of, on vegetable crops in Florida, 289–308; virus 1, natural history of, 306–307; virus 1, on Commelina nudiflora, 293; virus 1, on Geranium carolinianum, 293; wilt or yellows, taxonomy of Fusaria causing, 29
Cell stimulation, in plants, physiologic studies of pathogenic bacteria inducing, 31
Cephalocereus senilis, disease of, 9

Ceratostomella, pini, on pine attacked by Southern pine beetle, 7; sp., associated with disease of Platanus orientalis in Pennsylvania, 22; ulmi, air currents as possible carrier of, 442–443; ulmi, technique for isolation from bark beetles, 37–38; ulmi, transmission of, through root grafts, 1039–1040

Cercospora, arachadis, resistance of peanuts to, 971; beticola, control in southern Minnesota by dusting and spraying, 234–243; personata, resistance of peanuts to, 971

Cercosporella herpotrichoides, Wojnowicia graminis associated with, 406, 408

Cereal crops, winter, Wojnowicia graminis a parasite of, 405–415

Ceresan, use of, for control of seedling blight, of barley, 318–326

Cereus peruvianus, disease of, 9

Chaetochloa italica, Ustilago crameri on, in North China, 648–649

Chaetocnema, denticulata, vector of Aplanobacter stewarti, 32; pulicaria, vector of Aplanobacter stewarti, 32

Chalaropsis thielavioides, cause of root rot of Chinese elms, 653

Chamaecyparis lawsoniana, crown gall on, 894

CHANDLER, W. H., and HOAGLAND, D. R., Comments on article by ANTONI KOZLOWSKI on "Little leaf or rosette of fruit trees in California," 522

Charcoal, effect on infectivity of tobacco mosaic virus, 915

Chayota edulis, susceptible to celery virus 1, 392, 398

Cheiranthus, host for spotted wilt in California, 17

Chemical, splitting of the tomato "combination-streak" virus complex, 33; studies on the virus of tobacco mosaic IV. Some effects of different chemical agents on infectivity, 899–921; studies on the virus of tobacco mosaic V. Determination of optimum hydrogen-ion concentrations for purification by precipitation with lead acetate, 922–930

Chemicals, effect of, on sclerotia of Sclerotium rolfsii on strawberries, 966; effect on "combination streak" virus of tomatoes, 864–874

Chenopodium murale, susceptible to celery virus 1, 389, 395

Cherries, leaf spot caused by Coccomyces hiemalis, 132–151

Cherry, leaf spot, control by copper phosphate mixture, 32

CHESTER, K. STARR, The antigenicity of the plant viruses, 702–714; Serological evidence in plant-virus classification, 686–701; Serological evidence in the study of the relationships of certain plant viruses, 10

Chlidanthus fragrans, infection with Stagonospora curtisii, 264

Chloramine-T, inactivation of virus by, 706

Chloropicrin, control of root-knot nematode with, 67–90; effect of, on number of root-knot galls on pineapple, 72–74

CHRISTENSEN, J. J., see JOHNSON, I. J.; and E. C. STAKMAN, Relation of Fusarium and Helminthosporium in barley seed to seedling blight and yield, 309–327

CHRISTIE, J. R., book review of Plant parasitic nematodes, by T. Goodey, 287; the development of root-knot nematode galls, 10–11

CHUPP, CHARLES, Macrosporium and Colletotrichum rots of turnip roots, 269–274

Cineraria roots, Pythium sp., in, 14

Citrullus vulgaris, susceptible to celery virus 1, 381, 384, 398; susceptibility to Phymatotrichum omnivorum, 559–577

Citrus, areolate spot, caused by **Leptosphaeria bondari, n. sp.**, 884–886; factors influencing mycorrhizal associations in roots, 645–647; fungi on, nature of inhibition between, 763–775

CLARKE, A. E., see TOMPKINS, C. M.

CLAYTON, C. N., see ARMSTRONG, G. M.

CLAYTON, E. E., A new and important factor in the epidemiology of tobacco leaf diseases, 11; and J. A. STEVENSON, Nomenclature of the tobacco downy mildew fungus, 516–521

Clematis virginiana, inoculation experiments with Puccinia rubigo-vera, 663–671

Climate, cause of little leaf or rosette of fruit trees in California, 275–278
Coccomyces hiemalis, ascospores, 152; control by eradicant fungicide, 23; discontinuous variation in, 155, 156; effect of carbon from various sources on mycelium, 147, 148; effect of formalin on conidia, 139; effect of hydrogen-ion concentration on conidial germination, 138, 139; effect of nitrogen from various sources on mycelium growth, 145–147; effect of staling products on, 141, 142; growth of mycelium on various media, 140, 141; infection, 152, 153, 154; length and size of conidia from various hosts, 151, 152; variability in monosporic cultures, 26–27, 131–159
COCHRAN, L. C., see NELSON, RAY
Cocos plumosa, Penicillium vermoeseni on, in California, 896
COFFMAN, F. A., see HUMPHREY, H. B.
Cold resistance of oats, effect of crown rust on, 28
Coleoptile of barley, discoloration of, by Fusarium, 313–318; discoloration of, by Helminthosporium, 313–318
Coleosporium solidaginis on aster, control of, 17; varietal susceptibility to, 17–18
Colletotrichum, caulicolum, synonym of **C. truncatum**, n. comb., 122; fragariae, cause of anthracnose and wilt of strawberry, 973; gloesporioides, 886; higginsianum, cause of turnip rot, 272–274; higginsianum, spore size, 273; rot of turnip roots, 269–274; **truncatum**, n. comb., compared with C. lindemuthianum, 125; **truncatum**, n. comb., comparison with other Colletotrichum spp., 122–125; **truncatum**, n. comb., on eggplant in Mississippi, 967; **truncatum**, n. comb., on garden and Lima beans, 121–125
Colonial bent, leaf spot of, caused by **Helminthosporium erythrospilum**, n. sp., 345–361
Colonies of Coccomyces hiemalis, effect of temperature on diameter, 143, 144; variability of, 134
Commelina, communis, susceptible to celery virus 1, 389, 394; communis, transmission of Southern celery-mosaic virus to, 951; nudiflora, host of Southern celery-mosaic virus, 289–308, 377; nudiflora, spread of Southern celery-mosaic virus on, in Florida, 299–308; nudiflora, susceptible to celery virus 1, 389, 394; nudiflora, virus, mosaic, mechanical transmission of, to pineapple, 10; nudiflora, virus, mosaic, symptoms compared with yellow spot on pineapple, 10
Comparative studies of certain cultures of Puccinia rubigo-vera and P. tomipara on wild grasses, 657
Complement, -fixation reaction, more sensitive than precipitin reaction, 709; -fixation tests, for viruses, 691
Conglutination, see Mez reaction
Conidia, of Coccomyces hiemalis, length and size from various hosts, 151, 152; of Taphrina deformans, copulation, 56–57; of T. deformans, hibernation, 50–51
Conidial production, effect of ultraviolet light on, of Coccomyces hiemalis, 144, 145
Conifers, crown gall on, 894
CONNER, H. A., A. J. RIKER, and W. H. PETERSON, Biochemic studies on the metabolism of crown-gall and hairy-root bacteria and on the composition of crown galls, 11
Constitution of The American Phytopathological Society, 445
Control, of apple scab in relation to leaf growth rate, 373–375; of Cronartium species by Tuberculina maxima, 259, 260; of potash hunger and fusarium wilt in cotton, 969; of root knot, 67–90; of seedling blight of barley by seed treatment, 318–326; of Stewart's disease of corn, by treatment of seed with bacteriophage, 371–372; of Stilbum cinnabarinum on fig, 219, 220; of sugar-beet leaf spot in southern Minnesota, 234–243
Convolvulus, arvensis, not infected by Anguillulina dipsaci, 1006; arvensis, pathologico-anatomical changes in woodiness of fruit, 554; sp., susceptible to celery virus 1, 389, 396
COOK, H. T., Occurrence of oospores of Peronospora effusa with commercial

spinach seed, 11–12; and J. A. CALLENBACH, Spinach seed treatments in Virginia, 12

COOK, M. T., Relation of insect injuries and root diseases in sugar cane, 12

COOLEY, J. S., Relation of host vigor to apple infection with Xylaria mali, 12–13

COONS, G. H., and J. E. KOTILA, Influence of preceding crops on damping off of sugar beets, 13

Copper, absorption by conidia of Sclerotinia fructicola, 17; compounds, insoluble, as substitutes for Bordeaux, 40; effect of, on prevention of and recovery from tobacco frenching, 1075; fungicidal sprays containing, penetration of plant tissues by, 368–370; fungicides, as peach sprays, 972; -lime-arsenite preparations, fungicidal properties of, 23–24; oil-soluble, use of, as a fungicide, 368–370; phosphate mixture, promising fungicide, 32–33; resinate, oil-soluble, use of, as fungicide, 368–370; sulphate-lime, as dust for control of sugar-beet leaf spot, 234–243

Cordylanthus pilosus highly resistant to snapdragon rust, 985

Corn, bacterial leaf blight of, effect of, on invasion by Diplodia zeae, 1113; bacterial wilt, insect vectors of, 32; Diplodia zeae on, factors affecting infection with and spread of, in tissues, 1113–1114; Helminthosporium of, seed transmission of, 1109–1112; infection with Phytomonas vascularum, 997–998; loss from ear rots in the United States, 281–283; reduction in yield from smut, 223–233; seed, fungi in, 1111, 1112; seed, treatment of, with bacteriophage, for control of Stewart's disease, 371–372; smut, nature of variation, 34; smut, physiologic specialization, 34; Stewart's disease of, bacteriophage in relation to, 371–372; Stewart's wilt disease, host range, 992–1002; sweet, see sweet corn; sweet, spread of Southern celery-mosaic virus on, 289–308

Corticium vagum on potatoes under irrigation, 748–762

Coryphantha valida, disease of, 9

Cotton, angular leaf spot of, control of, 970, 972; angular leaf spot of, in Mississippi, 973; bacterial leaf spot, dissemination of organism, 971; fusarium wilt of, 975; fusarium wilt of, control of, 969; lint, effect of Phymatotrichum root rot on commercial grade of, 109–110; lint, effect of Phymatotrichum root rot on length of, 107–109; Phymatotrichum root rot, reduction in yield due to, 107–109; potash hunger, control of, 969; -root-rot fungus, effect of ammonia nitrogen on, 967–968; -root-rot fungus, Phymatotrichum omnivorum, parasitic on watermelon, Citrullus vulgaris, 559–577; root rot, Phymatotrichum, effect on convolution of fiber, 128–129; root rot Phymatotrichum, effect on ribbon width of fiber, 128; root rot, Phymatotrichum, effect on thickness of fiber, 128; seed, effect of Phymatotrichum root rot on quality of, 110–112; seed treatment, results from ten years' work, 972; seed, treatments in South Carolina, 970; seedlings, the etiology of damping off, 968–969; verticillium wilt disease of, 972; -wilt fungus, the reaction of several isolations of, to toxic dyes, 970

Cottons, wilt-resistant, adapted to the Gulf Coastal Plains, 974

Covered smut, an effective and easily applied method of inoculating barley seed with, 1038–1039

Cowpea, root-knot nematode on, 72–74; var. Black Eye, as test plant, infectivity of tobacco -ring spot and cucumber-mosaic viruses, 487–489

CRANDALL, B. S., see HARTLEY, CARL; see JACKSON, L. W. R.; see LAMBERT, E. B.

Crinum powelli, infection with Stagonospora curtisii, 264; S. curtisii on, 263

Cripple and dark stripe on pineapples in Hawaii, 25–26

Cronartium, cerebrum, parasitised by Tuberculina maxima, 255; cerebrum, T. maxima on aecia of, 1117–1118; coleosporioides, parasitised by T. maxima, 255–256; pyriforme, parasitised by T.

maxima, 255; ribicola, host of T. maxima, 255; ribicola, observations on T. maxima, a parasite of, 253–261; ribicola, spore stages of, dates of production in Pacific Northwest, 1104–1108

CROSIER, WILLARD, Joined spores of Ascochyta viciae, 283–284; and DONALD REDDICK, Some ecologic relations of Phytophthora infestans, 13

Crown, gall, biochemic studies, 11; gall, control, 6; gall, effect on apple trees, 975; gall, effect of, on apple trees in Orchard, 1099–1103; gall on conifers, 894; gall on red raspberry, 5–6; gall on Sequoia; gall, physiologic studies of bacteria, 31; rust, effect of, on cold resistance in oats, 28; rust, effect of, on composition of oats, 28–29; rust, effect of, on water economy of oats, 28–29; rust, effect of, on yield of oats, 28

Crucifers, symptoms of virus disease, 640; virus disease, 640–644

Cryptosporella umbrina, n. comb., cause of brown canker of rose, description of, 886–889

Cucumber, direct effect of Bordeaux mixture on early production of, 974; mosaic, inactivation of virus at different hydrogen-ion concentrations, 487–489; mosaic virus, similarity of Southern celery-mosaic to, 947

Cucumis, anguria, susceptible to celery virus 1, 389, 398; melo, susceptible to celery virus 1, 381, 398; sativus, susceptible to celery virus 1, 381, 382, 398

Cucurbita, maxima, susceptible to celery virus 1, 381, 398; pepo condensa, susceptible to celery virus 1, 381, 382, 398; pepo, Southern celery-mosaic virus on, 304–305; pepo, susceptible to celery virus 1, 381, 398

Cultural, studies on the virus of tobacco mosaic, 1035–1037; variations of Thielaviopsis basicola, 1011–1018

Culture, media, determination of pH of solid, 362–367; of Aplanobacter stewarti, for production of bacteriophage, 371

Cultures, from barley seed, 310; monosporic, varability in, of Coccomyces hiemalis, 131–159; of Puccinia rubigovera, 657; of P. tomipara, 657

Cupressus, spp., crown gall on, 894

Curly top, of beets, retention of virus by aphids, 175; on pansy, not transmitted by seed, 894; outbreak of, on pansy in Oregon, 894; (tomato yellows) in tomatoes, graft *versus* insect transmissions of, 844–853

CURRENCE, T. M., see LEACH, J. G.

Cuticle, blackberry, penetration by germinating aeciospores of Gymnoconia interstitialis, 453–474

Cyanogas, control of root-knot nematode with, 86–87

Cynoglossum amabile, susceptible to celery virus 1, 386, 396

Cytological study of the resistance of apple varieties to Gymnosporangium juniperi-virginianae, 30

Cytology of secondary spore formation in Ganoderma, 426–429

Dactylis glomerata, leaf smut on, 811; Ustilago clintoniana on, 815, 816

Damping off, control by sand culture of seedlings, 15; of alfalfa, caused by pythiaceous fungi, 8; of alfalfa on acid and neutral Iowa soils, 8–9; of alfalfa on acid soils, 8–9; of alfalfa on neutral soils, 8–9; of alfalfa, treatment of soil, 9; of flower seedlings, formaldehyde treatment, to control, 18; sugar beet, influence of preceding crops on, 13; the etiology of, of cotton seedlings, 968–969

DANA, B. F., and F. P. MCWHORTER, An outbreak of curly top on pansy, 894

Dark stripe and cripple of pineapples in Hawaii, 25–26

DARLING, H. M., J. G. LEACH, and F. A. KRANTZ, Scab resistance in potato seedlings, 13–14

DARROW, GEORGE M., Susceptibility of raspberry species and varieties to leaf spot (Mycosphaerella rubi) at Beltsville, Maryland, 961–962; Root rots and degeneration of the strawberry, review of papers on, 523–525

Dates of production of the different spore stages of Cronartium ribicola in the Pacific Northwest, 1104–1108

Datura, meteloides, susceptible to celery virus 1, 392, 397; stramonium, frenching of, 1070; stramonium, susceptible to celery virus 1, 398

Daucus carota, susceptible to celery virus 1, 381, 395

DAVEY, A. E., see LEACH, L. D.; and L. D. LEACH, Toxicity of compounds of ammonia to Sclerotium rolfsii, 895–896

DAVIDSON, ROSS W., see LAMB, HOWARD

DAVIS, W. H., Summary of investigations with Ustilago striaeformis parasitizing some common grasses, 810–817

DECKER, PHARES, see TAUBENHAUS, J. J.

Deficiency, mineral, of soil, effect of, on tobacco frenching, 1074–1081

Delphinium, consolida, susceptible to celery virus, 1, 386, 387, 395; host for spotted wilt in California, 17

Demonstration of plant-parastic nematodes in host tissues, 1026–1030

Dendroctonus frontalis, carrier of microorganisms in pine, 7

Dent corn, injury by fungi following seed-coat injury, 24; pathologic significance of seed-coat injury, 24; seed treatment, 24

Descriptive key for plant viruses, 328–343

Diabrotica duodecimpunctata vector of Aplanobacter stewarti, 32

Dianthus caryophyllus, bacterial leaf spot of, caused by Phytomonas woodsii, 114–120

Diaporthe, citri, effect of, on growth of Phytophthora spp. in culture, 766; citri, filtrate from, effect of, on growth of Phytophthora citrophthora, 769–772; eres, on rose petals in Louisiana, 888–889; umbrina, cause of brown canker of rose, transferred to **Cryptosporella umbrina**, n. comb., 887–889

Diffusible nature of the inhibitory agent produced by fungi, 1031–1034

Diplodia, natalensis, bark disease of Tahiti lime trees caused by, 966; root rot of strawberry caused by species of, 974–975; zeae, factors affecting infection with and spread of, 1113–1114; zeae, in corn seed, 1111

Diploidization of coalescing haploid pustules of Puccinia helianthi, 1087–1090

Dipotassium phosphate, effect on infectivity of tobacco-mosaic virus, 910

Disease, bark of Tahiti lime trees caused by Phomposis citri and Diplodia natalensis, 966; of peach tree recently discovered in California, 897; of sorghum, caused by a hybrid smut, 375–376; resistance, breeding peanuts for, 971–972; Stewart's, of corn, bacteriophage in relation to, 371–372

Diseases of cultivated plants, book review, 444

Disinfection, of barley seed, 310; tobacco bed soil, 972

Dissemination, of Southern celery-mosaic virus on vegetable crops in Florida, 289–308; of the bacterial leaf-spot organism, 971

Dissociation of Fusarium niveum in soil, 30–31

Distribution of intermediate types of barley smuts, 28

DIXON, LAWRENCE F., RUTH A. MCLEAN, and FREDERICK A. WOLF, Initiation of downy mildew of tobacco in North Carolina in 1934, 628–639

DOAK, K. D., and P. L. FISHER, Mycorrhizal and pseudomycorrhizal infections of pine roots during first year's growth, 14

Downy mildew, incidence on tobacco as related to seed-bed practices, 636–638; incidence on tobacco as related to seed beds on old and new sites, 630–636; incidence on tobacco from air-borne sporangia, 636; initiation on tobacco in North Carolina in 1934, 628–639; of tobacco, control, 19, 971; tobacco, nomenclature of fungus, 516–521

DRECHSLER, CHARLES, A leaf spot of bent grasses caused by **Helminthosporium erythrospilum**, n. sp., 344–361; Occurrence of a species of Aphanomyces on roots of spinach and flax, 14–15; A Pythium species of the megalacanthum

type in Cineraria roots and the relation of putrefaction to parasitism among the Pythiaceae, 14

DUGGAR, B. M., Thermal inactivation of some tobacco viruses: standardization, technique, and illustrative data, 15; and D. F. MCALLISTER, Some factors affecting "longevity" *in vitro* of viruses of tobacco mosaic and of tobacco ring spot, 15

DUNEGAN, J. C., Phytophthora disease of peach seedlings, 800–809; see ROBERTS, JOHN W.

DUNLAP, A. A., Sand culture of seedlings as a damping-off control, 15

Dust, copper sulphate-lime for control of sugar-beet leaf spot, 234–243; formaldehyde, and treatment of barley seed for seedling blight, 325–326

Dutch, disease, elm, transmission by root grafts, 1039–1040; elm-disease fungus, air currents as possible carrier of, 442–443

DYE, H. W., A neglected factor in the planning and interpretation of fungicidal tests, 15–16

Dyes, toxic to Fusarium vasinfectum, 971

DYKSTRA, T. P., A top-necrosis virus found in some apparently "healthy" potatoes, 1115–1116

EDDINS, A. H., Sclerotinia rot of Irish potatoes in Florida, 974; Soil treatment with sulphur and limestone for control of bacterial wilt of potatoes, 16

Effect, of carbon dioxide on the pH gradient of the sugar beet and on the feeding of the curly-top vector, 891; of certain chemicals on the "combination streak" virus of tomatoes, 864–874; of clean fallow on field populations of the root-knot nematode, Heterodera marioni, 891; of crown gall, hairy root, and woolly aphids on apple trees in the orchard, 1099–1103; of crown rust on the cold resistance of oats, 28; of crown rust on yield, water economy, and composition of oats, 28–29; of light intensity and temperature on infection of Hope wheat by Puccinia graminis tritici, 1041–1066; of light on the initiation of rust infection, 715–725; of mosaic on transpiration of the bean, 18; of proteoclastic enzymes on purified preparations of tobacco mosaic virus, 33

Eggplant, fruit rot of, in Mississippi, caused by **Colletotrichum truncatum**, n. comb., 967; fruit rot of, in Mississippi, caused by Gloeosporium melongena, 967

Eggplants, anthracnose of, 967

Electric soil sterilization, study of, 29–30

Electrodes, quinhydrone and antimony, for determining pH of solid culture media, 362–367

ELLIOTT, CHARLOTTE, see POOS, F. W.

Elm, Dutch disease, transmission by root grafting, 1039–1040

Elms, Chinese, root rot, 652–654

Elymus spp., inoculation experiments with Puccinia rubigo-vera, 663–671

Emilia, host for spotted wilt in California, 17; sagittata, susceptible to celery virus 1, 386, 388, 398

Endoxerosis of lemon fruits as affected by the application of different amounts of irrigation water, 897

Environment, effect of, on balance between host and parasite, 1059–1063; effect of, on infection by Puccinia graminis tritici, 1054–1055

Enzymes, proteoclastic, effect of, on tobacco-mosaic virus, 33

Epidemiology of tobacco leaf diseases, effect of driving rains, 11; effect of fertilizers, 11; effect of topping, 11

Erwinia, amylovora, filtrability of, 728–729; amylovora, longevity in honeybee environment, 20; amylovora, modes of entry into flowers, 20; amylovora, relation of carbohydrate nutrition of apple to infection by, 26; carotovora, filtrability of, 728–729

Etch, severe, tomato, serological evidence in classification of virus, 687–700

Euchlaena mexicana, infected by Aplanobacter stewarti, 32; susceptible to celery virus 1, 392, 393, 394

Evidence of the overwintering of Plasmopara australis in fruits of Sicyos angulatus, 27
Experiment to determine the susceptibility of flax to verticilliosis, 892
Experimental error in greenhouse pot tests with sugar beets, factors affecting, 1019–1025
Experiments, on physiologic specialization and nature of variation in Ustilago zeae, 34; on the control of the root-knot nematode in the field with chloropicrin and other chemicals, 67–90; on the multiplication of curly-top virus in the beet leaf hopper, 891
Explanation of Rhiomyza hypogaea, 898
EYER, J. R., Observations on the pathological histology and phyto-chemistry of psyllid yellows, 895
EZEKIEL, WALTER N., see STROMAN, G. N.; see TAUBENHAUS, J. J.; and J. J. TAUBENHAUS, Field trials of pentachlorethane, tetrachlorethane, and xylol as affecting phymatotrichum root rot and host plants, 16

Factors affecting infection with a spread of Diplodia zeae in host tissue, 1113–1114
Fagopyrum esculentum, susceptible to celery virus 1, 392, 393, 395
Fasciation of sweet peas: a bacterial disease, 36
FIFE, J. M., and V. L. FRAMPTON, The effect of carbon dioxide on the pH gradient of the sugar beet and of the feeding of the curly-top vector (title only), 891
Fig, Stilbum disease of, in Louisiana, 208–222
Filtrability, of bacteria, 728–729; of certain plant pathogenic bacteria, 728–729
Filtration, quantitative studies on, of tobacco-mosaic virus, 601–617
Fire-blight organism, longevity in honeybee environment, 20
Firmiana simplex, blight disease of, in Florida, 38
FISCHER, GEORGE W., Comparative studies of certain cultures of Puccinia rubigovera and P. tomipara on wild grasses, 657–685
FISHER, P. L., see DOAK, K. D.
Flax, experiment to determine susceptibility to verticilliosis, 892; roots, Aphanomyces sp. in, 14–15
Foeniculum vulgare, susceptible to celery virus 1, 381, 396
FORBES, I. L., see HART, HELEN
Formaldehyde, dust, and treatment of barley seed, for seedling blight, 325–326; hot, for potato seed treatment, 760; injury from, 18; treatment for damping off, 18
Formalin, effect on germination of conidia of Coccomyces hiemalis, 139
FOSTER, H. H., Infection of apple leaves by Physalospora cydoniae, 16–17
Foxtail, yellow, infection with Phytomonas stewarti, 997
Fragaria sp., black stele root rot, 5; Phytophthora on, 5
FRAMPTON, V. L., see FIFE, J. M.
Frankliniella sp., transmission of spotted-wilt virus by, 17
FREITAG, J. H., see SEVERIN, H. H. P.; and H. H. P. SEVERIN, Experiments on the multiplication of curly-top virus in the beet leaf hopper (title only), 891
FREMONT, THERESE, see REED, H. S.
Frenching, Nicotiana spp. susceptible to, 1070; plants susceptible to, 1070–1071; tobacco, see tobacco, frenching; tobacco, studies on, 1067–1084
Fructification in Macrosporium spp., method for inducing, 31–32
Fungi, inhibitory agent produced by diffusible nature of, 1031–1034; parasitic on citrus, nature of inhibition between, 763–775
Fungicidal, sprays containing copper, penetration of plant tissues by, 368–370; tests, planning and interpretation, 15–16
Fungicide, copper phosphate mixture, a promising, 32–33; red copper oxide, 21; studies on sulphur as, 35–36; use of oil-soluble copper as, 368–370; use of oil-soluble copper as, compared with Bordeaux, 368–370
Fungicides, Ceresan, for control of seedling blight, of barley, 318–326; copper, as peach sprays, 972; development of

eradicant, 23; effect on growth of apples as factor in comparative tests, 15–16; eradicant, control of certain diseases by, 23; formaldehyde dust, and treatment of barley seed, for seedling blight, 325–326; properties of certain copper-lime-arsenite preparations, 23–24
Fungous blight of barley seed, 310, 325
Fungus spot of azalea flowers, 38
Further, evidence of the seed-borne nature of Peronospora destructor, 35; studies on the longevity of sclerotia of Phymatotrichum omnivorum, 36
Fusaria causing celery wilt, taxonomy of, 29
Fusarium, apii, cause of wilt of celery, 29; apii, var. pallidum, cause of wilt of celery, 29; cactacearum, on cactus in Italy, 9; cacti maxonii, on cactus in Italy, 9; conglutinans, growth association, with other organisms, 9; culmorum, on winter wheat, 409; graminearum, predominant species in barley seed, 310; in barley seed, as affected by seed treatment, 318–326; in barley seed, prevalence of, by localities, 311–313; in barley seed, relation of, to discoloration of coleoptile, 313–318; in barley seed, relation of, to germination, 313–318; in barley seed, relation of to root rot, 313–318; in barley seed, relation of, to seedling blight, 309–327; in barley seed, relation of, to stand, 313–318; in barley seed, relation of, to stunting, 313–318; in barley seed, relation of, to yield, 309–327; lini, 29; lycopersici, growth association, with other organisms, 9; moniliforme, growth association, with other organisms, 9; moniliforme, in corn seed, 1111, 1112; niveum, 971; niveum, carried in seed of watermelons, 969; niveum, dissociation in soil, 30–31; pathogenicity of different strains to China aster, 734–738; sp., on winter wheat, 409; spp., associated with bottom rot of calla rhizomes, 39; spp., associated with damping off of cotton, 969; spp., section sporotrichiella, common in barley seed, 310; strains, in relations to wilt of China aster, 733–747; vasinfectum, carried in seed of okra, 979; vasinfectum, tested for reaction to toxic dyes, 970–971; wilt, control of, in cotton, 969; wilt, of cotton, 975; wilt, on celery, forms of, 29; wilt on muskmelon in Minnesota, 25; wilt, seeds of watermelons and okra as possible carriers of, 969; yellows of celery, 578–600
Gaillardia, host for spotted wilt in California, 17
Galanthus sp., infection with Stagonospora curtisii, 264
Gall, crown, effect on apple trees, 975
Ganoderma, secondary spore formation in, cytology of, 426–429
GARDNER, M. W., see TOMKINS, C. M.; C. M. TOMKINS, and O. C. WHIPPLE, Spotted wilt of truck crops and ornamental plants, 17
Garlic, nematode of, longevity in soils, 1009, 1010
Genotypes, of Neurospora sitophila, serically active carbohydrates of, 39; of N. tetrasperma, serically active carbohydrates of, 39
Geranium carolinianum, host of Southern celery-mosaic virus, 290–308; susceptible to celery virus 1, 389, 390, 395
German foxtail millet, infection with Phytomonas stewarti, 997
Germination, method of, for demonstrating Fusarium in barley seed, 313–315; method of, for demonstrating Helminthosporium in barley seed, 313–315; of conidia of Coccomyces hiemalis, effect of hydrogen ion and disinfectants, 138, 139; of urediospores of Puccinia graminis tritici, effect of plant extracts on, 1049–1050
Gibberella saubinetii, growth association, with other organisms, 9; in corn seed, 1111; on dent corn, 24; scab, on barley, recent outbreaks of, 309
Gilia capitata, susceptible to celery virus 1, 386, 396
GITMAN, L., and E. BOYCHENKO, Manual of diseases of new bast fiber plants (Russian), book review, 655
Gloeosporium, leaf spot, of orchids, 435–437; melongena, resemblance to, of

fungus from anthracnose of eggplant, 967

Glomerella, cingulata, comparison with G. rubicola, 93–100; cingulata, growth association with other organisms, 9; gossypii, losses caused by, to cotton seedlings, 968–969; rubicola, cause of white-bud disease of raspberry, 93; rubicola, comparison with G. cingulata, 93–100; rubicola, cultural characters, 93–95; rubicola, host relationships of, 92–93; rubicola, pathogenicity of, 96–97

Gloxinia, host for spotted wilt in California, 17

Glycine apios, Rhizoctonia sp. on, in Florida, 38

Godetia, host for spotted wilt in California, 17

GODFREY, G. H., Experiments on the control of the root-knot nematode in the field with chloropicrin and other chemicals, 67–90; The demonstration of plant-parasitic nematodes in host tissues, 1026–1030; and C. EMLEN SCOTT, New ecenomic hosts of the stem- and bulb-infesting nematode, 1003–1010

GOLDSWORTHY, M. C., see ROBERTS, JOHN W.; and E. L. GREEN, Availability of copper in Bordeaux mixture residues and its absorption by conidia of Sclerotinia fructicola, 17

GOODEY, T., Plant parasitic nematodes, book review, 287

Graft, knot, control by antiseptics and antiseptic tape, 203–206; *versus* insect transmissions of curly top in tomatoes (tomato yellows), 844–853

Grafts, root, transmitting Ceratostomella ulmi, 1039–1040

GRAHAM, T. W., Nuclear phenomena in Helminthosporium gramineum, 284–286; see VERRALL, A. F.

Grass, Sudan, infection with Phytomonas stewarti, 996–997

Grasses, bent, leaf spot of, caused by **Helminthosporium erythrospilum**, n. sp., 344–361; comparative studies of Puccinia on, 657–685

GRATZ, L. O., see KINCAID, R. R.

GREEN, E. L., see GOLDSWORTHY, M. C.; see ROBERTS, JOHN W.

Growth, association of microorganisms, 9; of Agaricus campestris on plots treated with sodium chlorate, 440–442; rate, in various media, of mycelium of Coccomyces hiemalis, 140, 141

GUTERMAN, C. E. F., Control of aster leaf rust, 17–18; see BURKHOLDER, WALTER H., and L. M. MASSEY, A liquid formaldehyde treatment to control damping off of flower seedlings, 18

Gymnoconia, interstitialis, infection by penetration of the cuticle, 453–474; juniperi-virginianae, cytological study of resistance of apple varieties to, 30; juniperi-virginianae, nature of resistance, of apple varieties to, 30

HAENSELER, C. M., see ALLEN, M. C.

HAFSTAD, G. E., see STAKMAN, E. C.

Hairy root, biochemic studies, 11; control by antiseptics and antiseptic tape, 192–207; effect of, on apple trees, 975; effect of, on apple trees in orchard, 1099–1103; of apple, caused by Phytomonas rhizogenes, 32; of apple, environmental factors influencing development of, 32

HAMILTON, JAMES M., see HORSFALL, JAMES G.

Haploid pustules, coalescing, in Puccinia helianthi, 1085–1090

Hardiness, relation of, to resistance to leaf spot on raspberries, 962

HARRIS, HUBERT A., Morphologic studies of Septoria lycopersici, 790–799

HARRISON, A. L., The effect of mosaic on transpiration of the bean, 18

HART, HELEN, and I. L. FORBES, The effect of light on the initiation of rust infection, 715–725; and KAROL ZALESKI, The effect of light intensity and temperature on infection of Hope wheat by Puccinia graminis tritici, 1041–1066

HARTLEY, CARL, and B. S. CRANDALL, Vascular disease in poplar and willow, 18–19

Heat treatment for the cure of yellows and rosette of peach, 24

HEDGCOCK, GEORGE G., Notes on the occurrence of Tuberculina maxima on the aecia of Cronartium cerebrum, 1117–1118

Helianthus annuus, infection with Puccinia helianthi, effect of light on, 715–725

Helminthosporium, effect of, on stem-rust infection, 1054; **erythrospilum,** n. sp., cause of leaf spot of bent grasses, 344–361; **erythrospilum,** n. sp., cause of leaf spot of bent grasses, as affected by temperature, 347–349; **erythrospilum,** n. sp., cause of leaf spot of bent grasses, cultural characters of, 354–359; **erythrospilum,** n. sp., cause of leaf spot of bent grasses, distribution of, 344–350; **erythrospilum,** n. sp., cause of leaf spot of bent grasses, effect of weather condition on, 347–349; **erythrospilum,** n. sp., cause of leaf spot of bent grasses, morphology of, 350–359; **erythrospilum,** n. sp., cause of leaf spot of bent grasses, pathological effects of, 345–349; **erythrospilum,** n. sp., cause of leaf spot of bent grasses, symptoms of, 345–349; **erythrospilum,** n. sp., cause of leaf spot of bent grasses, technical description of, 360; **erythrospilum,** n. sp., compared with Helminthosporium triseptatum, 349–350 gramineum, nuclear phenomena in, 284–286; in barley seed, as affected by seed treatment, 318–326; in barley seed, prevalence of, by localities, 311–313; in barley seed, relation of, to discoloration of coleoptile, 313–318; in barley seed, relation of, to germination, 313–318; in barley seed, relation of, to root rot, 313–318; in barley seed, relation of, to seedling blight, 309–327; in barley seed, relation of, to stand, 313–318; in barley seed, relation of, to stunting, 313–318; in barley seed, relation of, to yield, 309–327; inaequalis, growth association with other organisms, 9; sativum, growth association with other organisms, 9; sativum, inhibition of growth by a bacterium, 1032, 1033; sativum, predominant species in barley seed, 310; sp., in corn seed, 1111, 1112; triseptatum, compared with H. **erythrospilum,** n. sp., 349–350

HENDERSON, R. G., Control of downy mildew of tobacco, 19

Hendersonia, graminis, comparison with Wojnowicia graminis, 412; herpotrichia, comparison with Wojnowicia graminis, 412; hirta, comparison with Wojnowicia graminis, 412; secalina, comparison with Wojnowicia graminis, 412

HENRY, W. D., Studies on the soft-rot and colon-typhoid-dysentery groups of bacteria. 1. Pathogenicity, 19; see ORTON, C. R.

HENSON, LAWRENCE, Apothecium production in Sclerotinia trifoliorum and S. sclerotiorum, 19–20

Heterocaryosis in Helminthosporium gramineum, 285

Heterodera marioni, control experiments, with chloropicrin, 69–86; control experiments, with Cyanogas and potassium xanthate, 86–87; development in tomato seedlings, 10–11; effect of chloropicrin, on number of root galls, caused by, 72–74; soil fumigation, for control of, 69–86

Heterothallism in Peronospora parasitica, 8

HEWITT, J. LEE, To visualize a distinction between viruses and organisms, 892

HIGGINS, B. B., Breeding peanuts for disease resistance, 971–972; Results from ten years' work on cotton-seed treatment, 972

HILBORN, M. T., see STEINMETZ, F. H.

HILDEBRAND, E. M., Longevity of the fireblight organism in the honeybee environment, 20; see RIKER, A. J.; and L. H. MACDANIELS, Modes of entry of Erwinia amylovora into the flowers of the principal pome fruits, 20

HINES, LEE, see STAKMAN, E. C.

Hippeastrum, inoculations of Stagonospora on, 262, 263; vittatum, infection with Stagonospora curtsii, 264

HOAGLAND, D. R., see CHANDLER, W. H., 522

HOERNER, G. R., and D. C. SMITH, A new canker of hops in Oregon, 437–439

HOGGAN, ISME A., see JOHNSON, JAMES; and JAMES JOHNSON, A virus of crucifers and other hosts, 640–644

HOLBERT, J. R., P. E. HOPPE, and A. L. SMITH, Some factors affecting infection with and spread of Diplodia zeae in the host tissue, 1113–1114
Holcus, sorghum, infection with Phytomonas stewarti, 993–996; sorghum, infection with P. vascularum, 1000; sorghum, see sorghum; sorghum, susceptible to celery virus 1, 392, 394; sudanensis, infection with P. stewarti, 996–997; sudanensis see Sudan grass
HOLST E. C., see BRAMBLE, W. C.
HOLTON, C. S., Studies on seven differentiating characteristics of two physiologic forms of Tilletia tritici, 1091–1098
Honeybee, environment, longevity of fireblight organism in, 20
HOPPE, P. E., see HOLBERT, J. R.
Hops, a new canker of, in Oregon, 437–439
Hordeum, jubatum, reaction of, to stem rust infection, 1046; sp., susceptibility to hybrids between Ustilago hordei and U. medians, 5; sp., susceptibility, to U. hordei, 5; sp., susceptibility, to U. medians, 5; spp., inoculation experiments with Puccinia rubigo-vera, 663–671; vulgare, occurrence of U. medians, 28; vulgare, see barley; vulgare, U. hordei, albino strain, monosporidial lines, compatible with normal U. hordei and U. medians, 27–28
Hormones in malt extract, 147
Horse-radish, virus disease of, 640
HORSFALL, JAMES G., and JAMES M. HAMILTON, Some fungicidal possibilities of red copper oxide, 21
Host, not influencing virus precipitin reaction, 709; -parasite relationship in bean rust, 39; range of Phytomonas stewarti and P. vascularum, 992–1002; range of Southern celery-mosaic virus, 377–404; tissues, demonstration of plant-parasitic nematodes in, 1026–1030
Hosts, natural and differential, of plant viruses, 330–343; new economic, of stem- and bulb-infesting nematode, 1003–1010; of Southern celery-mosaic virus, 290–308; perennial, of Southern celery-mosaic virus, importance of, 307
HUBERT, ERNEST E., Observations on Tuberculina maxima, a parasite of Cronartium ribicola, 253–261
HUMPHREY, H. B., and F. A. COFFMAN, A portable rust-inoculation chamber, 279–281
Hunger, potash, control of, in cotton, 969
HUTCHINSON, W. G., Resistance of Pinus sylvestris to a gall-forming Peridermium, 819–843
Hybrid smut of sorghum, 375–376
Hybridization, between Sphacelotheca sorghi and Sorosporium reilianum, 375–376; between Ustilago hordei and U. medians, 5
Hydrogen-ion, concentration, effect of, on conidial germination of Coccomyces hiemalis, 138, 139; concentration, effect of, on inhibition between Diaporthe citri and Phytophthora species, 768; concentration, of soil, effect of, on tobacco frenching, 1073–1074; concentrations, determination of optimum for purification of tobacco-mosaic virus with lead acetate, 922–930; determination of, in solid culture media, 362–367; rates of inactivation of virus at different concentrations, 475–492
Hymenocallis, calathina, infection with Stagonospora curtisii, 264; littoralis, Phyllosticta hymenocallidis on, 263
Hystrix patula, inoculation experiments with Puccinia rubigo-vera, 663–671

Immunity, acquired, in Zinnia from cucumber mosaic, 776–789
Impatiens biflora, inoculation experiments with Puccinia rubigo-vera, 663–671
Importance of sanitation in controlling crown gall of the red raspberry, 5–6
Improved method for inducing spore fructification in certain species of Macrosporium, 31–32
Inactivation of virus and precipitin reactivity, 703–707
Incubation, of the virus of pea mosaic in the aphid, Macrosiphum gei, 31; period, of pea mosaic in aphid, 160–177
Infection, by Puccinia graminis tritici, effect of light intensity and temperature

on, 1041–1066; Gymnoconia interstitialis, penetration of cuticle by germinating aeciospores, 453–474; of fig by Stilbum cinnabarinum, 218; with Coccomyces hiemalis on various species of prunus, 152, 153, 154

Infectivity, effect of phosphate buffers on tobacco-mosaic virus, 618–627; of tobacco mosaic virus, effects of different chemical agents on, 899–921

Influence, of nutrition on host susceptibility of yellow tobacco mosaic, 33–34; of Phytomonas tumefaciens and P. rhizogenes on actual acidity of certain liquid and agar substrata, 854–863

Inheritance of resistance to common bean mosaic in field and garden beans, 875–883

Inhibition between certain fungi parasitic on citrus, nature of, 763–775

Inhibitory agent, produced by fungi, diffusible nature of, 1031–1034

Inoculation, effect of methods on virus of tobacco mosaic, 913; method, for rust transfer, 661; of seed barley with covered smut, an effective and easily applied method, 1038–1039; tests with Phytomonas stewarti and P. vasculara, 21

Inoculations, portable chamber for, 279–281

Insect, dissemination of azalea flower spot, 38; transmissions, *versus* graft of curly top in tomatoes (tomato yellows), 844–853; vectors of bacterial wilt of corn, 32; vectors of pea mosaic viruses, 31

Insecticide, nicotine, added to oil-soluble copper fungicide, 369

Insects, relation to root rot of sugar cane, 12; transmission of Southern celery-mosaic virus by, 290–308; vector of plant viruses, 330–343

Insoluble copper compounds as substitutes for Bordeaux, 40

Internal necrosis of bean seeds, 726–728

Interpretation of Oregon tip blight on a basis of causal viruses, 897–898

Intraspecific reactional differences of host plants to Puccinia, 683

Invalidity of genus Stagonosporopsis, 419

Ipomoea, batatas, susceptible to celery virus 1, 381, 384, 396; purpurea, susceptible to celery virus 1, 386, 396

Irrigation, effect of, on Rhizoctonia sclerotial development, 754–757

Isolate *vs.* isolant

IVANOFF, S. S., book review, Bolesti na kulturnite rastenija. (Diseases of cultivated plants) by D. ATANASOFF, 444; Inoculation tests with Phytomonas stewarti and P. vasculara, 21; Studies on the host range of Phytomonas stewarti and P. vascularum, 992–1002; see RIKER, A. J.; and A. J. RIKER, Resistance of sweet corn to bacterial wilt, 21–22

JACKSON, L. W. R., and B. S. CRANDALL, A phytophthora root and collar rot of Pinus resinosa seedlings, 22; and BAILEY SLEETH, A new disease affecting Platanus orientalis in the Eastern United States, 22

Japanese varnish tree, see Firmiana simplex

JENKINS, ANNA E., see BITANCORUT, A. A.; and LEWIS E. WEHMEYER, Transfer of Diaporthe umbrina to the genus Cryptosporella, 886–889

JOHANN, HELEN, see MACKIE, W. W.

JOHANNSON, H., see NEWTON, MARGARET

JOHNSON, E. M., An example of spread of veinbanding from potatoes to tobacco, 650–652; see VALLEAU, W. D.; and W. D. VALLEAU, Cultural variations of Thielaviopsis basicola, 1011–1018

JOHNSON, FRANK H., Cultural studies on the virus of tobacco mosaic, 1035–1037

JOHNSON, I. J., and J. J. CHRISTENSEN, Relation between number, size and location of smut infections to reduction in yield of corn, 223–233

JOHNSON, JAMES, book review, Virus diseases of plants. A bibliography by D. ATANASOFF, 444; see HOGGAN, ISME A., and ISME A. HOGGAN, A descriptive key for plant viruses, 328–343

JOHNSON, T., see NEWTON, MARGARET

JONES, L. R., see RIKER, REGINA S.

Juniperus spp., crown gall on, 894

KADOW, KENNETH J., The raspberry white-bud disease and its relation to bitter rot of apples, 91–103; and H. W. ANDERSON, The value of zinc sulphate as a peach spray ingredient, 22–23

KAUFERT, FRANK, The production of asexual spores by Pleurotus corticatus, 23

KEITT, G. W., Progress in the development of eradicant fungicides, 23; and D. H. PALMITER, Fungicidal properties of certain copper-lime-arsenite preparations, 23–24

KENT, G. C., see MELHUS, I. E.

Key, descriptive, for plant viruses, 328–343

KILMER, F. B., see RIKER, A. J.

KIMMEY, J. W., see MIELKE, J. L.

KINCAID, R. R., The effect of certain environmental factors on the germination of Florida cigar-wrapper tobacco seeds, 975; and L. O. GRATZ, Soil-temperature studies on Florida cigar-wrapper tobacco, 975

KOEHLER, BENJAMIN, Pathologic significance of seed-coat injury in dent corn, 24

Koeleria cristata, Wojnowicia graminis on, 408

KOTILA, J. E., see COONS, G. H.

KOZLOWSKI, ANTONI, Little leaf or rosette of fruit trees in California, 275–278; see CHANDLER, W. H., for comment upon former author's article on "Little leaf or rosette of fruit trees in California," 522

KRANTZ, F. A., see DARLING, H. M.

KUNKEL, L. O., Heat treatment for the cure of yellows and rosette of peach, 24

Laboratory and field studies on sulphur as a fungicide, 35–36

LAMB, HOWARD, ERNEST WRIGHT, and ROSS W. DAVIDSON, A root rot of Chinese elms, 652–654

LAMBERT, E. B., and B. S. CRANDALL, A phytophthora wilt of black-locust seedlings, 24–25

Late blight on tomatoes, 7; seed-borne, 7

Lathyrus odoratus, infected with Anguillulina dipsaci, 1006; infection with pea mosaic, 162, 163; see sweet pea

Layia, host for spotted wilt in California, 17

LEACH, J. G., see DARLING, H. M., and T. M. CURRENCE, Resistance to fusarium wilt in muskmelon, 25

LEACH, L. D., see DAVEY, A. E.; and A. E. DAVEY, Soil amendments for Southern sclerotium rot of sugar beets, 896; and A. E. DAVEY, Toxicity of low concentrations of ammonia to mycelium and sclerotia of Sclerotium rolfsii, 957–959

Lead acetate, determination of optimum hydrogen-ion concentrations with, 922–930

Leaf, rust, of aster, control, 17; rust, of aster, varietal susceptibility to, 17–18; scorch, of narcissus, due to Stagonospora curtisii, 262, 263; spot, angular, of cotton in Mississippi, 973; spot, bacterial, dissemination of organism, 971; spot, bacterial, of carnation, 114–120; spot, of bent grasses, caused by **Helminthosporium erythrospilum**, n. sp., 344–361; spot, of bent grasses, caused by **H. erythrospilum**, n. sp., compared with spot due to H. triseptatum, 349–350; spot, of orchids, 435–437; spot, of sugar beet, control of, in southern Minnesota, 234–243; spot, red, due to Stagonospora curtisii, 262, 263; spot, susceptibility of raspberries to, 961–962

Leaves, apple-spur, growth rate of, in relation to scab, 373–375; apple-spur, rate of increase in area of, 373–375

LE CLERG, E. L., Dusting and spraying experiments for the control of sugar-beet leaf spot in southern Minnesota, 234–243; Factors affecting experimental error in greenhouse pot tests with sugar beets, 1019–1025

Lemon, endoxerosis of fruits as affected by the application of different amounts of irrigation water, 897

Leptosphaeria, bondari, n. sp., cause of areolate spot of citrus, 884–886; **bondari**, n. sp., technical description of, 886; citricola, 886

Lesions, necrotic, relation to localization of tobacco-mosaic viruses in Nicotiana, 968

Lespedeza sericea, stem blight of, 976

Leucojum vernum, infection with Stagonospora curtisii, 264

LEVINE, M. N., book review, Manual of diseases of new bast fiber plants (Russian), by L. GITMAN and E. BOYCHENKO, 655;

see STAKMAN, E. C.; and E. C. STAKMAN, Uniform rust nurseries indicate decreasing severity of stem rust, 25
LI, H. W., see TU, CHIH
Libocedrus decurrens, crown gall on, 894
Life history of Taphrina deformans, 41–66
Light, effect of, on initiation of rust infection, 715–725; intensity, effect of, on infection of Hope wheat by Puccinia graminis tritici, 1041–1066; intensity, effect of reduced, on infection by P. graminis tritici, 1055–1058; ultraviolet, effect on conidial production of Coccomyces hiemalis, 144, 145
Ligniera junci, root parasite of Stellaria media, observations on, 893
Lilium, auratum, susceptible to celery virus 1, 386, 394; longiflorum, susceptible to celery virus 1, 386, 387, 394
Lime bark disease of Tahiti lime trees caused by Phomopsis citri and Diplodia natalensis, 966
Linaria, six species immune from snapdragon rust, 985
LINFORD, M. B., Cripple and dark stripe of pineapples in Hawaii, 25–26
LINK, GEORGE K. K., Relation of carbohydrate-nitrogen nutrition to disposition of apple to infection by Erwinia amylovora, 26; Suitability of the Chicago soil-nutrient temperature tank for phytopathologic studies, 26; see WILCOX, HAZEL W.
Little leaf, of fruit trees in California, 275–278; rosette, California fruit trees, 522
Longevity *in vitro* of plant viruses, 330–343
Losses, corn-ear rots in the United States, 281–283; from corn smut infections, 222–233
Lychnis viscaria, susceptible to celery virus 1, 386, 395
Lycopersicum esculentum, frenching of, 1070; late blight of, 7; see tomato, susceptible to celery virus 1, 381, 383, 397; vehicle of immunization antigen, 689
Lycoris squamigera, infection with Stagonospora curtisii, 264
MACDANIELS, L. H., see HILDEBRAND, E. M.
MACKIE, W. W., Aeroplane dusting with sulphur to combat stem rust of wheat, 892–893; HELEN JOHANN, and N. E. STEVENS, An unidentified species of Sphaeropsis on maize in California, 893
Macrosiphum, gei, carrier of pea mosaic, 174; gei, incubation of virus of pea mosaic in, 31; gei, transmission of virus of pea mosaic by, 31; pisi, incubation of virus of pea mosaic, in, 31; pisi, incubation period of pea mosaic in, 160–177; pisi, transmission of virus of pea mosaic in, 31
Macrosporium, blight on tomato, control of, 968; herculeum, cause of, rot on turnips, 270; inducing spore fructification in certain species of, 31–32; rot of turnip roots, 269–274; seed treatment for, 272
MAGIE, ROBERT O., Variability of monosporic cultures of Coccomyces hiemalis, 26–27, 131–159
MAINS, E. B., Rust resistance in Antirrhinum, 977–991
Maize, spread of Southern celery-mosaic virus on, 289–308; unidentified species of Sphaeropsis on, in California, 893
Malt extract, effect on making amino acids available in growth of conidia of Coccomyces hiemalis, 147
Malus sp., bitter rot of, relation to raspberry white-bud disease, 91–112; Physalospora cydoniae, variation in isolates, 16–17; trunk canker, caused by Phytophthora cactorum, 5; trunk canker, control, 5; trunk canker, symptoms, 5; Xylaria mali, 12
MANDENBERG, E. C., see BANFIELD, W. M.
MANEY, T. J., see PLAGGE, H. H.
MASSEY, L. M., see GUTERMAN, C. E. F.
MCALISTER, D. F., see DUGGAR, B. M.
MCCLINTOCK, J. A., see SHERBAKOFF, C. D.
MCLEAN, RUTH A., see DIXON, LAWRENCE F.
MCWHORTER, FRANK P., The properties and interpretations of tulip-breaking viruses, 898; The symptoms of narcissus mosaic developed within the plant, 896–897; see DANA, B. F.; and J. A. MILBRATH, The interpretation of Oregon tip blight on a basis of causal viruses, 897–898

Mechanical transmission of two viruses to pineapple, 10
Media, Lima-bean agar used for monosporic cultures, Coccomyces hiemalis, 133
Medicago sativa, damping off, 8; see alfalfa
Megalonectria, nigrescens, 217; pseudotrichia, ascigerous stage of Stilbum cinnabarinum, 215–217; pseudotrichia, on fig in Louisiana, 215–217
MEHRLICH, F. P., Nonsterile soil leachate stimulating to zoosporangia production by Phytophthora sp., 432–435
MEIER, F. C., Microorganisms in the atmosphere of Arctic regions, 27
MELHUS, I. E., and G. C. KENT, Evidence of the overwintering of Plasmopara australis in fruits of Sicyos angulatus, 27; and C. S. REDDY, A practical method of organization for longtime phytopathological field experiments (title only), 27
Mercuric chloride, antiseptic in control of hairy root, graft knot, crown gall, 196–207; for potato seed treatment, 760
Mercury, organic, treatment for Rhizoctonia of potato, 760
Mesembryanthemum crystallinum, susceptible to celery virus 1, 386, 388, 395
Method, for demonstrating Fusarium in barley seed, by germination, 313–315; for demonstrating Helminthosporium in barley seed, by germination, 313–315; method of determining the rate of decay in wood, 35; of inoculating seed barley with covered smut, 1038–1039; of obtaining bacteriophage, from cultures of Aplanobacter stewarti, 371; photoelectric, for measurement of leaf area, 374
Methods for determining pH of solid culture media, 362–367
MEZ reaction with plant-virus materials, 691–692
MICHAILOWA, P. V., Pathologico-anatomical changes in the tomato incident to development of woodiness of the fruit, 539–558
Microflora of barley seed, 310–313
Microorganisms, growth association of, 9; in the atmosphere of Arctic regions, 27; infecting pines attacked by Dendroctonus frontalis, 7; pathogenic on cactus, 9
Microsiphum solanifolii, vector of potato veinbanding virus, 650
MIELKE, J. L., and J. W. KIMMEY, Dates of production of the different spore stages of Cronartium ribicola in the Pacific Northwest, 1104–1108
MILBRATH, J. A., see MCWHORTER, FRANK P.
Mildew, downy, initiation on tobacco in North Carolina in 1934, 628–639; downy, of tobacco, value of sprays for control of, 971
MILES, L. E., Angular leaf spot of cotton in Mississippi in 1934, 973; Report of one year's test of copper fungicides and other new materials as peach sprays, 972; The verticillium wilt disease of cotton, 972
Millet, breeding for smut resistance in North China, 648–649; common, infection with Phytomonas stewarti, 997; German foxtail, infection with P. stewarti, 997
Mineral deficiency not direct cause of tobacco frenching, 1074–1081
Mites on potatoes, 654
MIX, A. J., The life history of Taphrina deformans, 41–66
Mixed cultures of fungi associated with citrus fruits, 763–775
Moisture, soil, effect of, on development of celery yellows, 585–590
Monilia sp., cause of little leaf or rosette of fruit trees in California, 275–278
Monochaetia mali, cause of apple canker, 1116–1117
Monosporidial lines of albino strain compatible with normal Ustilago hordei and U. medians, 27–28
Monosporium cactacearum, on cactus in Italy, 9
MOORE, M. B., and C. C. ALLISON, An albino strain of barley smut, 27–28; and C. C. ALLISON, The distribution of intermediate types of barley smuts, 28
MOORE, W. D., see ANDRUS, C. F.
Morphologic studies of Septoria lycopersici, 790–799
Mosaic, aucuba, of potato, serological evidence in classification of virus, 687–700;

bean, effect on transpiration, 18; burn, certain strains of tobacco mosaic causing, 967; celery disease in California, 891; celery, Southern, classification of virus, 947–954; common bean, inheritance of resistance to, in field and garden beans, 875–883; cucumber, acquired immunity in Zinnia from, 776–789; cucumber, inactivation of virus at different hydrogen-ion concentrations, 487–489; cucumber, serological evidence in classification of virus, 687–700; cucumber, virus, similarity of Southern celery-mosaic to, 947; incubation period of pea, in aphid, 160–177; latent, of potato, serological evidence in classification of virus, 687–700; mild, of potato, serological evidence in classification of virus, 687–700; of cabbage, 6; of cabbage, effect on seed production, 6; of cabbage, symptoms, 6; of cabbage, transmission, 6; on roses, symptoms of, 8; on roses, transmission of, 8; on roses, varieties of, 8; pea, on Lathyrus odoratus, 162; pea, on Pisum sativum, 162; pea, on Trifolium incarnatum, 162; pea, serological evidence in classification of virus, 687–700; pea, transmitted mechanically, 175; quantitative studies on filtration of virus from tobacco, 601–617; retention of virus by pea aphid, 170–174; rose, yellow variety of, 8; Southern celery, causal factors in increase of, 301–304; Southern celery, dissemination of, on vegetable crops in Florida, 289–308; Southern celery, factors influencing field dissemination of, 304–306; Southern celery, host range of, 377–404; Southern celery, insect vector of, in Florida, 290–308; Southern celery, natural history of, 306–307; Southern celery, spread of, in seed beds, 302; Southern celery, wild hosts of, in Florida, 290–308; sugar-cane, severe type of, 36–37; sugar-cane, symptoms, 36–37; symptoms of, developed within narcissus plant, 896–897; tobacco, causing mosaic burn, 967; tobacco, chemical studies of the virus of, 899–921; tobacco, chemical studies on virus, 475–492; tobacco, cultural studies on the virus of, 1035–1037; tobacco, determination of optimum hydrogen-ion concentrations for purification of virus by precipitation with lead acetate, 922–930; tobacco, effect of tannic acid on infectivity of virus, 931–937; tobacco, longevity of virus, factors affecting, 15; tobacco, necrotic lesions in Nicotiana, 968; tobacco, rates of inactivation of virus at different hydrogen-ion concentrations, 475–492; tobacco, virus, strains as affected by the nitrogen supply, 179–188; tobacco, yellow, influence of phosphorus and potassium on susceptibility to, 493–502; yellow tobacco, influence of nutrition on susceptibility to, 33–34; yellow tobacco virus, effect of nitrogen supply on susceptibility of tobacco, 184–188

Mottle, of potato, serological evidence in classification of virus, 687–700

Mottle, of potato, serological evidence in the cold resistance of oats, 28; Effect of crown rust on yield, water economy, and composition of oats, 28–29

Musa, cavendishii, susceptible to celery virus 1, 392, 394; sapientum, susceptible to celery virus 1, 392, 394

Muskmelon, resistance to fusarium wilt, 25

Mustard, virus disease of, 640

Mycelium, toxicity of low concentrations of ammonia to, of Sclerotium rolfsii, 957–959

Mycorrhiza, pine, 14

Mycorrhizal, and pseudomycorrhizal infections of pine roots during first year's growth, 14; associations, factors influencing formation and development in citrus roots, 645–647

Mycosphaerella, fragariae, mode of action of Bordeaux on, 969; rubi, susceptibility of raspberries to, 961–962; rubina, control by eradicant fungicide, 23

Myrothecium roridum, cause of black crown rot of greenhouse snapdragons in Texas, 969

Myzus persicae, vector of virus of crucifers, 640

Narcissus, inoculations of Stagonospora curtisii on, 262; leaf scorch of due to S. curtisii, 262, 263; mosaic symptoms de-

veloped within the plant, 896–897; sp., infection with S. curtisii, 264

NEAL, D. C., Further studies on the effect of ammonia nitrogen on growth of the cotton-root-rot fungus, Phymatotrichum omnivorum, in field and laboratory experiments, 967–968; Wilt-resistant cottons adapted to the Gulf Coastal Plains, 974

Necrosis, internal, of bean seeds, 726–728

Necrotic spotting, factor for, in Nicotiana tabacum, Mendelian dominant, 968

Neglected factor in the planning and interpretation of fungicidal tests, 15–16

NELSON, RAY, and L. C. COCHRAN, Taxonomy of the Fusaria that cause celery wilt (yellows), 29; Three forms of the fusarium wilt of celery, 29

Nematode, galls, caused by Heterodera marioni, 10; galls, development in tomato seedling, 10–11; root-knot, Heterodera marioni, effect of clean fallow on field populations of, 891; stem- and bulb-infesting, new hosts of, 1003–1010

Nematodes, associated with damping off of cotton, 969; location of, in diseased tissues, 1007; plant-parasitic, book review, 287; plant-parasitic, demonstration in host tissues, 1026–1030

Neomamillaria gubrowiana, disease of, 9

Neurospora, sitophila, serically active carbohydrates of genotypes of, 39; terasperma, serically active carbohydrates of, genotypes of, 39

New, blight of pea in California, 891–892; disease affecting Platanus orientalis in the eastern United States, 22; variety of Tilletia tritici in Montana, 40

NEWHALL, A. G., A study of electric soil sterilization, 29–30; see STUART, W. W.

NEWTON, MARGARET, H. JOHANNSON, and T. JOHNSON, A study of the carotenoid pigments of urediospores of wheat stem rust and four of its color variants, 30

Nicandra physalodes, susceptible to celery virus 1, 392, 397

Nicotiana, glauca, host for spotted wilt in California, 17; glutinosa, effect of mosaic burn on, 967; glutinosa, experimental material, tobacco virus inactivation tests, 477; glutinosa, nitrogen supply as affecting virus infection, 178–191; glutinosa, susceptible to celery virus 1, 392, 397; glutinosa, test of virus on, 901; glutinosa, tests for virus infectivity on, 923; necrotic lesions of tobacco-mosaic viruses in, 968; sanderae, susceptible to celery virus 1, 392, 397; sp., vehicle of immunization antigen, 689; spp., attempt to localize tobacco-mosaic virus on, 968; spp., susceptible to frenching, 1070; sylvestris, susceptible to celery virus, 1, 392, 397; tabacum, downy mildew, control of, 19; tabacum, mosaic causing spotting in, 968; tabacum, mosaic longevity of virus, 15; tabacum, mosaic virus, particle size of, 36; tabacum, ring spot, longevity of virus, 15; tabacum, ring spot, recovery and immunity from, 37; tabacum, see tobacco; tabacum, susceptible to celery virus 1, 391, 392, 397; tabacum, test of virus on, 901, 922; tabacum, var. ambalema, resistance of, to different viruses, 37; tabacum, var. Turkish, experimental material, influence of host nutrition on virus susceptibility, 494; tabacum, var. Turkish, experimental material, tobacco virus, inactivation tests, 477; tabacum, var. Turkish, nitrogen supply as affecting virus infection, 178–191; tabacum, viruses, thermal inactivation of, 15; tabacum, wildfire, on, in Pennsylvania, 6; tabacum, yellow mosaic, influence of nutrition on susceptibility, to, 33–34

NIEVES, RAIMUNDO, Experimental infection of rye, var. Petkus, by wheat bunt; Tilletia tritici and T. levis (in Spanish, English summary), 503–515; Infección experimental del centeno de Petkus (Secale cereale v. vulgare) por los caries del trigo: Tilletia tritici y T. levis (in Spanish with English summary), 503–515

Nitrogen, effect of, from various sources, on mycelium growth of Coccomyces hiemalis, 145–147; effect on host susceptibility to virus infection, 178–191

NOLAN, R. E., Sclerotium rolfsii on strawberries and the effect of certain chemicals on the sclerotia, 966; A root rot of straw-

berry caused by a species of Diplodia, 974–975
Nomenclature, downy mildew, tobacco, 516–521
Nonspecific reactions, eliminated by testing virus in one host with serum prepared from another, 709
Notes on the occurrence of Tuberculina maxima on the aecia of Cronatrium cerebrum, 1117–1118
Nuclei in Helminthosporium gramineum, 284–286
NUSBAUM, C. J., A cytological study of the resistance of apple varieties to Gymnosporangium juniperi-virginianae, 30
Nutrients, mineral, effect of deficiency of, on tobacco frenching, 1074–1081
Nutrition, carbohydrate-nitrogen, relation of, to infection of apple by Erwinia amylovora, 26; influence of, on host susceptibility to yellow tobacco mosaic, 33–34

Oak, Pleurotus corticatus on, 23
Oats, effect of crown rust on cold resistance of, 28; effect of crown rust on yield, water economy, and composition of, 28–29; winter, Wojnowicia graminis on, 408
Observations, on a species of Ligniera, a root parasite of Stellaria media, 893; on the pathological histology and phytochemistry of psyllid yellows, 895
Oil, copper resinate dissolved in, use of, as fungicide, 368–370
Okra, seeds of, as possible carriers of fusarium wilt, 969
Oleander, tubercle, physiologic studies of bacteria, 31
Olive, tubercle, physiologic studies of bacteria, 31
OLIVEIRA, J. M., The effect of clean fallow on field populations of the root-knot nematode Heterodera marioni (title only), 891
ONG, E. R., DE, The use of oil-soluble copper as a fungicide, 368–370
Onion seed, transmission of mildew by, 35
Orchids, Gloeosporium leaf spot of, 435–437
Ornamental plants infected by celery virus, 385
ORTON, C. R., Dissociation of Fusarium niveum in soil, 30–31; and W. D. HENRY, An internal necrosis of bean seeds, 726–728
OSBORN, H. T., Incubation of the virus of pea mosaic in the aphid, Macrosiphum gei, 31; Incubation period of pea mosaic in the aphid, Macrosiphum pisi, 160–177
Outbreak of curly top on pansy, 894

PADY, S. M., Aeciospore infection in Gymnoconia interstitialis by penetration of the cuticle, 453–474
PALMITER, D. H., see KEITT, G. W.; see RIKER, A. J.
Palms, ornamental, relation of Penicillium vermoeseni to a disease of, 896
Palustrex, copper resinate dissolved in, use of, as fungicide, 369
Pancratium maritimum, infection with Stagonospora curtisii, 264
Panicum miliaceum, infection with Phytomonas stewarti, 997
Pansy, curly top outbreak on, in Oregon, 894
Papain, effect of, on tobacco-mosaic virus, 33
Papaver, host for spotted wilt in California, 17
Parasitic, nematodes, plants, demonstration in host tissues, 1026–1030; studies, on tobacco frenching, 1071–1072
PARKER, HORACE, see BARRETT, J. T.
Parsley infected with Anguillulina dipsaci, 1005–1006
Particle size of three strains of tobacco-mosaic virus, 36
Pastinaca sativa, susceptible to celery virus 1, 381, 396
Pathogenicity, of monoconidial cultures of Fusarium causing celery yellows, 593–595; of soft-rot and colon-typhoid-dysentery groups of bacteria, 19; of Stilbum cinnabarinum on fig, 217, 218
Pathologic significance of seed-coat injury in dent corn, 24
Pathologico-anatomical changes, in tomato, incident to woodiness of fruit, 539–558; in woodiness of fruit of bindweed, 554

Pea, effect on retention in aphid of mosaic virus of immune plants, 173; mosaic, incubation in aphid, 31; mosaic, incubation of virus in aphid, 31; mosaic, incubation period in aphid, 160–177; mosaic, on Lathyrus odoratus, 162; mosaic, transmission by aphid, 31; new blight of, in California, 891–892

Peach, leaf curl, life history of organism, 41–66; leaf curl, occurrence of late infection, 49–50; leaf curl, origin of saprophytic phase, 63–64; leaf curl, spraying against, 48; phony disease, 24; rosette, heat treatment cure, 24; seedlings, phytophthora disease of, in Arkansas, 800–809; spray, zinc sulphate as an ingredient, 22–23; sprays, copper fungicides, etc., 972; tree disease recently discovered in California, 897; yellows, heat treatment cure, 24

Peanuts, breeding for disease resistance, 971–972

Pear, infection of, by Stilbum cinnabarinum, 219; leaf blight, control by copper phosphate mixture, 32

Pelargonium hortorum, susceptible to celery virus 1, 386, 395

Penicillium, oxalicum, on dent corn, 24; spp., inhibitory effect on other fungi, 9; vermoeseni, relation of, to a disease of ornamental palms, 896

Pentachlorethane, effect on phymatotrichum root rot and host plants, 16

Pentstemon, host for spotted wilt in California, 17

Peppers, spread of Southern celery-mosaic virus on, in Florida, 289–308

Pepsin, effect of, on tobacco-mosaic virus, 33

Peptone, bacteriologic, poor in growth hormones, 147

Perennial hosts of Southern celery-mosaic virus, importance of, 307

Peridermium, gall-forming, resistance of Pinus sylvestris to, 819–843

Peronospora, destructor, evidence for seed transmission, 35; effusa, occurrence of oospores with spinach seed, 11–12; hyoscyami, not cause of downy mildew of tobacco, 516;521; nicotianae, not cause of downy mildew of tobacco, in the United States, 516–521; parasitica, heterothallic strains, 8; parasitica, homothallic strains, 8; parasitica, in storage cabbage, 955–957; parasitica, oospore production, 8; parasitica, overwintering of, 957; tabacina, cause of downy mildew of tobacco, 516–521

PERSONS, T. D., Anthracnose disease of eggplants, 967

Pestalozzia sp., on Pinus resinosa seedlings, 22

PETERSON, W. H., see CONNER, H. A.

Petroleum oil, copper resinate dissolved in, use of, as fungicide, 369

Petroselinum hortense, infected with Anguillulina dipsaci, 1005, 1006; susceptible to celery virus 1, 381, 396

Petunia hybrida, frenching of, 1070; susceptible to celery virus 1, 385, 386, 398

pH, effect on precipitin reactivity of virus, 705

Phacelia, tanacetifolia, susceptible to celery virus 1, 386, 396; whitlava, susceptible to celery virus 1, 386, 396

Phaseolus, lunatus, **Colletotrichum truncatum**, n. comb., on, in southern and eastern United States, 121–125; vulgaris, **Colletotrichum truncatum**, n. comb., on, 121–125; vulgaris, infection with Uromyces appendiculatus, effect of light on, 715–725; vulgaris, mosaic, 18; vulgaris, nitrogen supply as affecting virus infection, 178–191; vulgaris, Tarsonemus latus on, 654; vulgaris, test of virus on, 901; vulgaris, tests for virus infectivity on, 923; vulgaris, var. Early Golden Cluster, experimental material, tobacco virus-inactivation tests, 477

Phenol, antiseptic tests in control of hairy root and crown gall, 194–202

Phleum pratense, physiologic form of Ustilago striaeformis on, 816

Phlox drummondii, susceptible to celery virus 1, 386, 388, 396

Phoenix, canariensis, Penicillium vermoeseni on, in California, 896; dactylifera, P. vermoeseni on, in California, 896

Pholidota imbricata, Gloeosporium leaf spot of, 435–437

Phoma, associated with bottom rot of calla rhizomes, 39; -like organism, associated with stem blight of Lespedeza sericea, 976
Phomopsis, citri, bark disease of Tahiti lime trees caused by, 966; on rose, 889
Phosphate dipotassium, effect on infectivity of tobacco-mosaic virus, 910
Phosphorus, nutrition, influence on host susceptibility to tobacco mosaic, 493–502
Photoelectric device, use of, for measuring increase in area of apple leaves, 373–374
Phyllosticta, disciformis, 886; hespidearum, 886; hymenocallidis, on Hymenocallis littoralis, 263; hymenocallidis, synonym of Stagonospora curtisii, 266; richardiae, associated with bottom rot of calla rhizomes, 39
Phymatotrichum, omnivorum, effect of ammonia nitrogen on growth of, 967–968; omnivorum, effect of Trichoderma lignorum on, 573–574; omnivorum, longevity of sclerotia, 36; omnivorum, parasitic on watermelon, 559–577; omnivorum, penetration studies on watermelon, 568–573; omnivorum, root rot, field trials with chemicals, 16; root rot, control with sulphur, 36; root rot, of cotton, effect on commercial grade of lint, 109–110; root rot, of cotton, effect on convolution of fiber, 128–129; root rot, of cotton, effect on length of lint, 107–109; root rot, of cotton, effect on quality of seed, 110–112; root rot, of cotton, effect on ribbon width of fiber, 128; root rot, of cotton, effect on thickness of fiber, 128; root rot of cotton, reduction in yield caused by, 107–109
Physalis, alkekengi, susceptible to celery virus 1, 381, 397; angulata, susceptible to celery virus 1, 389, 397; lagascae, susceptible to celery virus 1, 389, 397; peruviana, susceptible to celery virus 1, 392, 397; sp., susceptible to celery virus 1, 389, 397; virginiana, Bacterium tabacum on, 6
Physalospora cydoniae, infection of apple leaves by, 16–17; on apple, varietal susceptibility to, 17; variation in isolates, 16–17
Physiologic, forms, of Puccinia graminis tritici, prevalence, 34; forms, of Tilletia tritici, differentiating characteristics of, 1091–1098; specialization in Ustilago zeae, 34; studies of several pathogenic bacteria that induce cell stimulation in plants, 31; studies on tobacco frenching, 1072–1082
Physiology of soft-rot and colon-typhoid-dysentery groups of bacteria, 34–35
Phytolacca, americana, susceptible to celery virus 1, 389, 395; rigida, susceptible to celery virus 1, 389, 391, 395
Phytomonas, pruni, bacteriophage, particle diameter of certain plant viruses and, 938–946; rhizogenes, environmental factors influencing development of, 32; rhizogenes, influence of, on acidity of liquid and agar substrata, 854–863; stewarti, host range of, 992–1002; stewarti, inoculation tests, 21; stewarti, on different hosts, 21; stewarti, resistance of sweet corn to, 21; stewarti, see Aplanobacter; tumefaciens, influence of, on acidity of liquid and agar substrata, 854–863; vascularum, host range of, 992–1002; vascularum, inoculation tests, 21; woodsii, causal organism of bacterial leaf spot of carnation, 118–119; woodsii, pathogenicity of, 115–118
Phytophthora, associated with black stele root rot of strawberry, 5; cactorum, cause of trunk canker of apple, 5; cactorum, control of, on apple, 5; cactorum, on peach seedlings, in Arkansas, 800–809; cactorum, varietal susceptibility of apple to, 5; cinnamomi, 22; citrophthora, effect of Diaporthe citri, on growth of, in culture, 766; citrophthora, effect of Diaporthe citri, on production of reproductive structures of, in culture, 769; citrophthora, effect of filtrate from Diaporthe citri on growth of, 769–772; cryptogea, cause of root rot of China aster, in California, 895; cryptogea, first record of occurrence in America, 895; disease of peach seedlings, 800–809; infestans, dissemination, 13; infestans, effect of temperature and moisture, 13; infestans, germination of sporangia, 13;

infestans, on tomatoes, seed-borne, 7; megasperma, cause of root rot of cauliflower in California, 893–894; parasitica, effect of Diaporthe citri on growth of, in culture, 766; parasitica, reaction of, with Phytophthora citrophthora and Diaporthe citri, 768; parasitica, var. nicotianae, soil-temperature studies on, 975; root and collar rot of Pinus resinosa seedlings, 22; root and collar rot of P. resinosa seedlings, fungi associated with, 22; trunk canker of apple, 5; wilt of black-locust seedlings, 24–25; wilt of black-locust seedlings in Virginia, 24; zoosporangia production in nonsterile soil leachate stimulating to, 432–435

PIERCE, L., see ROBERTS, J. W.

PIERCE, W. H., The inheritance of resistance to common bean mosaic in field and garden beans, 875–883

PINCKARD, J. A., Physiologic studies of several pathogenic bacteria that induce cell stimulation in plants, 31

Pine, attacked by Dendroctonus frontalis, microorganisms infecting, 7; occurrence of Ceratostomella pini, a yeast and unnamed basidiomycete, 7; roots, mycorrhizal and pseudomycorrhizal infection, 14; -tar oil, copper resinate dissolved in, use of, as fungicide, 369

Pineapple, green spotting, relation to symbiont of Pseudococcus brevipes, 10; Heterodera marioni on, 67; root-knot nematode, effect of chloropicrin on number of root-galls, 72–74; root-knot nematode, soil fumigation for control of, 67–90; toxic effect on, of feeding by P. brevipes, 10; virus, Commelina nudiflora, symptoms compared with yellow spot, 10; virus, mosaic, mechanical transmission from C. nudiflora, 10

Pineapples, cripple and dark stripe in Hawaii, 25–26

Pinus, echinata, mycorrhizal and pseudomycorrhizal infection, 14; monticola, host of Cronartium species, 255; resinosa, mycorrhizal and pseudomycorrhizal infection, 14; resinosa, seedlings, phytophthora root and collar rot of, 22; strobus, mycorrhizal and pseudomycorrhizal infection, 14; sylvestris, resistance to gall-forming Peridermium, 819–843; taeda, mycorrhizal and pseudomycorrhizal infection, 14

PIRONE, P. P., An improved method for inducing spore fructification in certain species of Macrosporium, 31–32

Pisum sativum, infection with pea mosaic, 162

PLAGGE, H. H., and T. J. MANEY, Soggy breakdown of winter banana apples, 730–731

PLAKIDAS, A. G., Mode of action of Bordeaux on Mycosphaerella fragariae, 970

Plant, parasitic nematodes, book review, 287; parasitic nematodes in host tissues, demonstration of, 1026–1030; viruses, bases for description and identification of, 328–343; viruses, chief groups of, 332; viruses, descriptive key for, 328–343

Plants, miscellaneous, susceptible to celery virus, 391

Plasmopara australis, overwintering of, in fruits of Sicyos angulatus, 27

Platanus, orientalis, Ceratostomella sp. associated with new diseases of, 22; orientalis, new disease on, in Pennsylvania, 22; sp., bacteria associated with discoloration in, 19

Pleurotus corticatus, production of asexual spores, 23

Plowrightia morbosa, control by eradicant fungicide, 23

Poa, annua, leaf smut on, 812; annua, physiologic form of Ustilago striaeformis on, 816; pratensis, leaf smut on, 812; pratensis, physiologic form of U. striaeformis on, 816; sandbergii, Wojnowicia graminis on, 408

Polyporus, calcuttensis, secondary spore formation in, 426; ochroleucus, secondary spore formation in, 426

POOS, F. W., and CHARLOTTE ELLIOT, Bacterial wilt of corn and its insect vectors, 32

POPE, O. A., see YOUNG, V. H.

Poplar, vascular disease, 18–19

Population trends of physiologic form of Puccinia graminis tritici, 1930 to 1934, 34

Populus, deltoides, vascular disease, 18–19; nigra italica, vascular disease, 18–19

POST, THELMA B., see WEISS, FREEMAN
Potash hunger, control of, in cotton, 969
Potassium, nutrition, influence on host susceptibility to tobacco mosaic, 493–502; permanganate, inactivation of virus by, 706; xanthate, control of root-knot nematode with, 86–87
Potato, late blight, important factors in spread of, 13; nature of scab resistance, 13; psyllid yellows, observations on pathological histology and phyto-chemistry of, 895; reaction of seedlings of inbred lines to scab, 13–14; rhizoctonosis of, 748–762; seed treatments as control of rhizoctonosis, 758–760; scab, control with sulphur, 36; top-necrosis virus in apparently healthy, 1115–1116
Potatoes, bacterial wilt, soil treatment for control of, 16; example of spread of veinbanding to tobacco, 650–652; Irish, sclerotinia rot of, 974; mites on, 654
Precipitin, reaction of plant viruses, not affected by host, 709; reactivity, of plant viruses, effect of inactivation on, 703–707; tests, for viruses, 689–691
PRICE, W. C., Acquired immunity from cucumber mosaic in zinnia, 776–789; Classification of Southern celery-mosaic virus, 947–954
Primula, host for spotted wilt in California, 17
Production of asexual spores by Pleurotus corticatus, 23
Properties and interpretations of tulip-breaking viruses, 898
Prunus, cerasifera, as host of Coccomyces hiemalis, 151; cerasifera, infection trials with C. hiemalis, 154–155; cerasus, as host of C. hiemalis, 151; domestica, infection trials with C. hiemalis, 154–155; insititia, infection trials with C. hiemalis, 154, 155; mahaleb, as host of C. hiemalis, 151; munsoniana, infection trials with C. hiemalis, 154–155; pennsylvanica, as host of C. hiemalis, 151; persica, see peach; susceptibility of various species to C. hiemalis, 152
Pseudococcus, brevipes, effect of feeding on pineapple leaves, 10; brevipes, symbionts in relation to phytotoxis, 10
Pseudomonas tumefaciens, on conifers, 894; producing crown gall on Sequoia, 439
Pseudomycorrhiza, pine, 14
Pseudopeziza ribis, optimum pH for germination, 7; optimum temperature for germination, 7; pathogenicity, 7; variation in conidia, 7
Pseudotsuga taxifolia, crown gall on, 894
Puccinia, antirrhini, apparently heteroecious, 978; antirrhini, effect of light on infection by, 715–725; antirrhini, germination of urediospores at various temperatures, 979; antirrhini, hosts for aecial stage unknown, 978; antirrhini, occurrence in United States, Canada, Bermuda, England, Belgium, and Germany, 977; antirrhini, resistance of Antirrhinum to, 977–991; antirrhini, technic of inoculation, 979; coronata, effect of light on infection by, 715–725; coronata, effect of, on cold resistance in oats, 28; coronata, effect of, on composition of oats, 28–29; coronata, effect of, on water economy of oats, 28–29; coronata, effect of, on yield of oats, 28; graminis, color variant of, 30; graminis tritici, bacterial parasite of, 1053; graminis tritici, effect of light on infection by, 715–725; graminis tritici, infection by, effect of environment on, 1054–1055; graminis tritici, infection by, effect of high temperature on, 1058–1059; graminis tritici, infection by, effect of light intensity and temperature on, 1041–1066; graminis tritici, infection by, effect of light intensity, reduced on, 1055–1058; graminis tritici, infection by, effect of other organisms on, 1053–1054; graminis tritici, inoculation of Hope wheat with, by hypodermic syringe, 1050–1053; graminis tritici, population trends of physiologic forms, 34; graminis tritici, urediospore germination of, effect of plant extracts on, 1049–1050; graminis, urediospores, carotenoid pigments of, 30; helianthi, effect of light on infection by, 715–725; helianthi, haploid pustules coalescing in, 1085–1090; klebahni, synonym of Cronartium ribicola, 253; rubigo-vera, comparative morphology of urediospores and teliospores from differ-

ent cultures of, 671–674; rubigo-vera, comparative studies on wild grasses, 657–685; rubigo-vera, cultural experiments with, 663–674; sorghi, effect of light on infection by, 715–725; tomipara, comparative studies on wild grasses, 657–685; tomipara, cultural experiments with, 674–679; triticina, effect of light on infection by, 715–725

Pumpkin, sclerotinia rot in Montana, 39–40

Pycnidial formation in Septoria lycopersici, 792

Pycnospore, formation and germination of, in Septoria lycopersici, 796

Pycnospores, fused, of Ascochyta viciae, 283–284

Pyrus malus, Tarsonemus latus on, 654

Pythium, anandrum, behavior toward bacteria, 14; antagonistic action of Trichoderma on, 244–252; de baryanum, toxicity of filtrate of Trichoderma on, 251; mastophorum, behavior toward bacteria, 14; polymastum, behavior toward bacteria, 14; sp., in Cineraria roots, 14; spp., relation of putrefaction to parasitism, 14; ultimum, cause of poor germination of cotton seed, 969

Quinhydrone, electrode, use of, 362–367

RAMSEY, G. B., Peronospora in storage cabbage, 955–957

Ranunculus acer, inoculation experiments with Puccinia rubigo-vera, 667

Raspberries, black, possible source of streak infection in, 959–961

Raspberry, black, Gymnoconia interstitialis on, aeciospore infection by, 458; black, physiologic studies of bacterium causing overgrowths on canes of, 31; pathogenicity of Glomerella rubicola to, 96–97; red, crown gall on, control of, 5–6; susceptibility to leaf spot, 961–962; white-bud disease, caused by Glomerella rubicola, 93; white-bud disease, economic importance, 93; white-bud disease, host relationship, 92–93; white-bud disease, relation to bitter rot of apples, 93–100; white-bud disease, symptoms, 91–92

Rate of increase in area of apple-spur leaves, 373–375

Red, clover, crown rot in Kentucky, 19; copper oxide, fungicidal possibilities, 21; gum, Pleurotus corticatus on, 23

REDDICK, DONALD, Mites on potatoes, 654: see CROSIER, WILLARD

REDDY, C. S., see MELHUS, I. E.

Redtop, leaf spot of, caused by **Helminthosporium erythrospilum**, n. sp., 344–361

REED, H. S., and THERESE FREMONT, Factors that influence the formation and development of mycorrhizal associations in citrus roots, 645–647; and H. H. THORNBERRY, A peach-tree disease recently discovered in California, 897

Relation, of Fusarium and Helminthosporium in barley seed to seedling blight and yield, 309–327; of Penicillium vermoeseni to a disease of ornamental palms, 896

Report, of the annual meeting of the Southern Division of The American Phytopathological Society, 965; of the nineteenth annual meeting of the Pacific Division of The American Phytopathological Society, 890–891; of Summer Meeting, 451–452; of the twenty-sixth annual meeting of The American Phytopathological Society, 528–538

Resistance, disease, breeding peanuts for, 971–972; inheritance of, to common bean mosaic in field and garden beans, 875–883; of ambalema tobacco to different viruses, 37; of Antirrhinum to rust, 977–991; of apple varieties to Gymnosporangium juniperi-virginianae, cytological study of, 30; of celery to yellows, 590–593; of Pinus sylvestris to a gall-forming Peridermium, 819–843; of sweet corn to bacterial wilt, 21–22; to fusarium wilt in muskmelon, 25; to rust, in Antirrhinum, 977–991; to snapdragon rust, a simple dominant character, 988

Revision of constitution of The American Phytopathological Society, 445

Rhiomyxa hypogoea, explanation of, 898

Rhizoctonia, aerial, on beans in Florida, 38; antagonistic action of Trichoderma

on, 244–252; bud rot of strawberry, 965–966; causing damping off of cotton, not important, 969; solani, associated with bottom rot of calla rhizomes, 39; solani, causing bud rot of strawberry, 966; sp. in Florida, hosts for, 38; sp., on Pinus resinosa seedlings, 22

Rhizoctonosis, of potatoes, 758–762; relation of soil temperature to potato stem infection, 752–753

Ribes, Pseudopeziza ribis on, 6–7

RIKER, A. J., see CONNER, H. A.; see IVANOFF, S. S.; S. S. IVANOFF, and F. B. KILMER, Antiseptic solutions and antiseptic adhesive tape in relation to control of hairy root, crown gall, and other overgrowths on nursery apple trees, 192–217; D. H. PALMITER, and E. M. HILDEBRAND, Some environmental factors influencing the development of apple hairy-root and of Phytomonas rhizogenes, 32

RIKER, REGINA S., and L. R. JONES, Fusarium strains in relation to wilt of China aster, 733–747

Ring spot, of potato, serological evidence in classification of virus, 687–700; tobacco, inactivation of virus at different hydrogen-ion concentrations, 487–489; tobacco, longevity of virus, factors affecting, 15; tobacco, recovery and immunity from, 37; tobacco, serological evidence in classification of virus, 687–700

ROBERTS, JOHN W., An apple canker caused by Monochaetia mali, 1116–1117; L. PIERCE, M. A. SMITH, J. C. DUNEGAN, E. L. GREEN, and M. C. GOLDSWORTHY, Copper phosphate mixture: A promising fungicide, 32–33

ROLFS, F. M., Dissemination of the bacterial leaf spot organism, 971

Root, grafts, transmitting Ceratostomella ulmi, 1039–1040; knot, control experiments, with carbon bisulphide, 76–86; knot, control experiments, with chloropicrin, 69–86; knot, control experiments, with Cyanogas and potassium xanthate, 86–87; knot, effect of chloropicrin, on number of root galls, 72–74; -knot experiments, control of, 67–90; -knot nematode galls, caused by Heterodera marioni, 10; -knot, nematode galls, development in tomato seedling, 10–11; -knot nematode, on cowpea, 72–74; -knot nematode, see Heterodera marioni; knot, soil fumigation for control of, 69–86; rot, black, of apple, 976; rot, of aster, caused by Phytophthora cryptogea, 895; rot, of barley, as related to Fusarium in seed, 313–318; rot, of barley, as related to Helminthosporium in seed, 313–318; rot, of Chinese elms, 652–654; rot, of strawberry, caused by species of Diplodia, 974–975; rot, phymatotrichum, 16; rot, phymatotrichum, of cotton, effect on convolution of fiber, 128–129; rot, phymatotrichum, of cotton, effect on ribbon width of fiber, 128; rot, phymatotrichum, of cotton, effect on thickness of fiber, 128; rot, phymatotrichum, on watermelon, 559–577; rots and degeneration of the strawberry, book review, 523

Roots, factors influencing formation and development of mycorrhizal associations in citrus, 645–647

Rose, brown canker caused by **Cryptosporella umbrina, n. comb.**, 887–889; Diaporthe eres, on petals, in Louisiana, 888–889; mosaic, 8; mosaic, symptoms, of, 8; mosaic, types of, 8; mosaic, yellow variety of, 8; phomopsis on, 889; streak, symptoms, 8; streak transmission of virus, 8; streak, virus disease of, 7–8

Rosette, (Little leaf) California fruit trees, 522; peach, heat treatment for cure of, 24

ROSS, A. F., The effect of proteoclastic enzymes on purified preparations of tobacco-mosaic virus, 33

Rot, black crown, of greenhouse snapdragons caused by Myrothecium roridum, 969; foot, see foot rot; of China aster root, caused by Phytophthora cryptogea, 895; of turnips from Macrosporium and Colletotrichum, 269–274; phytophthora, of cauliflower root, 893–894; rhizoctonia bud, of strawberry, 965–966; root, of barley, as related to Fusarium in seed, 313–318; root, of barley, as related to Helminthosporium in seed, 313–318; root, see root rot; sclerotinia, of Irish

potatoes in Florida, 974; Southern sclerotium, of sugar beets, soil amendments for, 896; stem, sweet potato, control of, 976

Rots, loss from corn-ear, in the United States, 281–283

Rubus sp., crown gall, 5–6; white-bud disease, caused by Glomerella rubicola, 93; white-bud disease, economic importance, 93; white-bud disease, host relationship, 92–93; white-bud disease, symptoms, 91–92

RUDOLPH, B. A., An experiment to determine the susceptibility of flax to verticilliosis, 892; Report of the nineteenth annual meeting of the Pacific Division of The American Phytopathological Society, 890–891

Rust, Antirrhinum, resistance to, 977–991; bean, host-parasite relationship, 39; infection, effect of light on initiation of, 715; resistance, in Antirrhinum, 977–991; stem, of wheat, aeroplane dusting with sulphur to combat, 892–893

Rusts, portable inoculation chamber, 279–281

Rye, var. Petkus, experimental infection, by wheat bunt, 503–515; Wojnowicia graminis on, 408

RYKER, T. C., Fusarium yellows of celery, 578–600

Saccharum officinarum, inoculations with Phytomonas stewarti negative, 997; mosaic, 36–37; root disease, relation of insects to, 12; see sugar cane

Salicylic acid, antiseptic tests in control of hairy root and crown gall, 194–198

Salix caprea, vascular disease, 18–19

Salsify infected with Anguillulina dipsaci, 1003–1005

Salvia, host for spotted wilt in California, 17

Sanitation in control of crown gall, 5–6

Scab, of apple, leaf growth rate in relation to control of, 373–375; on barley, caused by Gibberella saubinetii, recent outbreaks of, 309; potato, nature of resistance to, 13; reaction of potato seedlings of inbred lines to, 13–14

Scabiosa atropurpurea, susceptible to celery virus 1, 386–398

SCHAAL, L. A., Rhizoctonosis of potatoes grown under irrigation, 748–762

Sclerotia, of Phymatotrichum omnivorum, longevity of, 36; of Sclerotium rolfsii, effect of certain chemicals on, 966; toxicity of low concentrations of ammonia to, of S. rolfsii, 957–959

Sclerotinia, americana, growth association with other organisms, 9; fructicola, absorption of Bordeaux mixture by conidia of, 17; fructicola, control by eradicant fungicide, 23; rot, of Irish potatoes in Florida, 974; rot, on pumpkin and squash, 39–40; sclerotiorum, cause of fruit rot of pumpkin and squash in Montana, 40; sclerotiorum, cause of potato rot in Florida, 974; sclerotiorum, on Canada thistle, 903; sclerotiorum, production of apothecium, 19–20; trifoliorum, production of apothecium, 19–20; wilt of Canada thistle, 963–964

Sclerotium, rolfsii, on strawberries and effect of certain chemicals on sclerotia of, 966; rolfsii, resistance of peanuts to, 972; rolfsii, soil treatment for, 896; rolfsii, toxicity of compounds of ammonia to, 895–896; rolfsii, toxicity of low concentrations of ammonia to mycelium and sclerotia of, 957–959; sp., on Pinus resinosa seedlings, 22

SCOTT, C. EMLEN, see GODFREY, G. H.

Seaside bent, leaf spot of, caused by **Helminthosporium erythrospilum**, n. sp., 345–361

Secale cereale, susceptible to celery virus 1, 393, 394

Seed, barley, blight of, 309–326; barley, cultures from 310; barley, disinfection of, 310; barley, method of inoculating with covered smut, 1038–1039; barley, microflora of, 310–313; barley, relation of Fusarium in, to discoloration of coleoptile, 313–318; barley, relation of Fusarium in, to germination of, 313–318; barley, relation of Fusarium in, to root rot, 313–318; barley, relation of Fusarium in, to seedling blight, 309–327; barley, relation of Fusarium in, to stand,

313–318; barley, relation of Fusarium in, to stunting, 313–318; barley, relation of Fusarium in, to yield, 309–327; barley, relation of Helminthosporium in, to discoloration of coleoptile, 313–318; barley, relation of Helminthosporium in, to germination, 313–318; barley, relation of Helminthosoporium in, to root rot, 313–318; barley, relation of Helminthosporium in, to seedling blight, 309–327; barley, relation of Helminthosporium in, to stand, 313–318; barley, relation of Helminthosporium in, to stunting, 313–318; barley, relation of Helminthosporium in, to yield, 309–327; barley, treatment of, for Fusarium, 318–326; barley, treatment of, for Helminthosporium, 318–326; beds, spread of Southern celery mosaic in, 302; blight, bacterial, of barley, 310–325; blight, fungous, of barley, 310–325; blight, of barley, caused by Alternaria, 310; -coat, injury in dent corn, pathologic significance of, 24; corn, treatment of, with bacteriophage, for control of Stewart's disease, 371–372; of cotton, effect of phymatotrichum root rot on quality of, 105–112; transmission of bacteria causing sweet pea fasciation, 36; transmission, of Bacterium malvacearum, 971; transmission, of Fusarium niveum, 969; transmission, of Fusarium vasinfectum, 969; transmission, of Helminthosporium of corn, 1109–1112; transmission, of Peronospora destructor, 35; transmission, of Plasmopara australis, 27; transmission, of spinach downy mildew, 11–12; transmission, of tomato late blight, 7; treatment, cotton, results from ten years' work, 972; treatment, of dent corn, 24; treatment of spinach, 12

Seedling, blight, of barley, effect of seed treatment on, 318–326; disease, of alfalfa, related to soil acidity, 421–425

Seeds, tobacco, effect of certain environmental factors on germination of Florida cigar-wrapper, 975

Segregation in barley smut, for chlamydospore-wall characters, 5; for head type, 5; for pathogenicity, 5; for sex, 5; in F_1, 5; in F_2, 5

Semesan B., treatment for rhizoctonia of potato, 760

Septoria lycopersici, morphologic studies of, 790–799

Sequoia, crown gall on, 439–440; spp., crown gall on, 894

Serically active (haptenic) carbohydrates of genotypes of Neurospora tetrasperma and N. sitophila, 39

Serological, evidence in plant-virus classification, 686–701; evidence in study of the relationships of certain plant viruses, 10; relationships of plant viruses, 10

Serology in virus classification, 688–701

Serum, preparation of immune, 688–689

Setaria, glauca, infection with Phytomonas stewarti, 997; glauca, inoculation with P. stewarti, 21; italica, var. Stramineofructa, infection with P. stewarti, 997

Severe type of mosaic on a sugar cane variety, 36–37

SEVERIN, H. H. P., see FREITAG, J. H.; and J. H. FREITAG, California celery mosaic disease, 891

SHAPOVALOV, MICHAEL, Chemical splitting of the tomato "combination-streak" virus complex, 33; Effect of certain chemicals on the "combination-streak" virus of tomatoes, 864–874; Graft *versus* insect transmission of curly top in tomatoes (tomato yellows), 844–853

SHARVELLE, E. G., see STAKMAN, E. C.

SHEAR, G. M., The growth of Agaricus campestris on plots treated with sodium chlorate, 440–442

SHERBAKOFF, C. D., and J. A. MCCLINTOCK, Effect of crown gall, hairy root, and woolly aphids on apple trees in the orchard, 975–976, 1099–1103; and G. M. STONE, Field studies of fusarium wilt of cotton, 975

SHUMWAY, CLYDE P., see TYLER, LEON J.

Sicyos angulatus, overwintering of Plasmopora angulatus in fruits of, 27

Silver, nitrate, inactivation of virus by, 706; possible compound with virus, 707

SLEETH, BAILEY, see JACKSON, L. W. R.

SMITH, A. L., see HOLBERT, J. R.

SMITH, C. O., Crown gall on conifers, 894; Crown gall on the Sequoia, 439–440; Inoculations of Stagonospora curtisii on

the Amaryllidaceae in California, 262–268

SMITH, D. C., see HOERNER, G. R.

SMITH, M. A., see ROBERTS, JOHN W.

SMUCKER, S. J., Air currents as a possible carrier of Ceratostomella ulmi, 442–443

Smut, covered, an effective and easily applied method of inoculating seed barley with, 1038–1039; covered kernel, and head smut, of sorghum, hybridization between, 375–376; millet, breeding for resistance in North China, 648–649; relation between infections and yield on corn, 223–233

Snapdragon, resistance to rust, 977–991; resistance to rust of commercial strains and varieties, 981–985; rust, resistance to, 977–991; rust-resistant selections, 982–985

Snapdragons, greenhouse, black crown rot of, in Texas caused by Myrothecium roridum, 969

SNYDER, WILLIAM C., A new blight of pea in California, 891–892

Sodium chlorate, growth of Agaricus campestris on plots treated with, 440–442

Soft, -rot bacteria, pathogenicity of, 19; -rot bacteria, physiology of, 34–35; scald, type of soggy breakdown in apples, 731

Soggy breakdown of winter banana apples, 730–731

Soil, acidity, relation to a seedling disease of alfalfa, 421–425; aeration, effect of, on tobacco frenching, 1073; amendments for Southern sclerotium rot of sugar beets, 896; disinfection, tobacco bed, 972; fumigation, for control of root knot, 69–86; leachate, nonsterile, stimulating to zoosporangia production in Phytophthora, 432–435; mineral deficiency of, effect of, on tobacco frenching, 1074–1081; reaction, effect of, on tobacco frenching, 1073–1074; sterilization, control of various microorganisms by electric, 29–30; sterilization, electric, 29; temperature in relation to Rhizoctonosis of potato, 752–754; -temperature studies on Florida cigar-wrapper tobacco, 975; texture, effect of, on tobacco frenching, 1072–1073; treatment with sulphur for control of potato bacterial wilt, 16

Solanum, aviculare, susceptible to celery virus 1, 392, 396; melongena, susceptible to celery virus 1, 381, 383, 397; nigrum, susceptible to celery virus 1, 389, 396; tuberosum, bacterial wilt, soil treatment for control, 16; tuberosum, nature of scab resistance in, 13; tuberosum, reaction of seedling of inbred lines to scab, 13–14; tuberosum, vehicle of immunization antigen, 689

Some, environmental factors influencing the development of apple hairy-root and of Phytomonas rhizogenes, 32; factors affecting infection with and spread of Diplodia zeae in the host tissue, 1113–1114

SORAUER, PAUL, Handbuch der Pflanzenkrankheiten, book review. Volume I, Part II, sixth edition, 525–527

Sorghum, hybrid smut of, 375–376; infection with Phytomonas stewarti, 993–996; infection with P. vascularum, 1000; inoculation with P. stewarti, 21; inoculation with P. vascularum, 21; smut, control with sulphur, 36

Sorosporium reilianum, and Sphacelotheca sorghi, hybridization between, 375–376

Southern celery, mosaic, spread of, in seed beds, 302; -mosaic virus, causal factors in increase of, 301–304; -mosaic virus, dissemination of, on vegetable crops in Florida, 289–308; -mosaic virus, factors influencing field dissemination of, 304–306; -mosaic virus, insect vector of, in Florida, 290–308; -mosaic virus, natural history of, 306–307; -mosaic virus, wild hosts of, in Florida, 290–308

SPENCER, ERNEST L., Effect of nitrogen supply on host susceptibility to virus infection, 178–191; Influence of nutrition on host susceptibility to yellow tobacco mosaic, 33–34; Influence of phosphorus and potassium on host susceptibility to yellow tobacco mosaic infection, 493–502; Studies on frenching of tobacco, 1067–1084

Sphaceloth eca sorghi, and Sorosporium reilianum, hybridization between, 375–376; control with sulphur, 35–36
Sphaeropsis, ambigua, 893; malorum, 893; sp., on Pinus resinosa seedlings, 22; unidentified species on maize in California, 893
Sphaerostilbe nigrescens, 217
Spinach, roots, Aphanomyces sp. in, 14–15; seed, occurrence of oospores of Peronospora effusa, 11–12; seed, treatment with various compounds, 12
Spinacia oleracea, see spinach; seed, occurrence of oospores of Peronospora effusa, 11–12; seed treatment with various compounds, 12; susceptible to celery virus 1, 381, 395
Spore formation, secondary, in Ganoderma, 426–429; secondary, in Polyporus calcuttensis, 426; secondary, in P. ochroleucus, 426; secondary, in Trametes lactinea, 426
Spores in air currents, 27
Sporotrichium, cactorium on cactus in Italy, 9; traversianum, on cactus in Italy 9
Spot, leaf, of bent grasses, caused by **Helminthosporium erythrospilum,** n. sp., 344–361; leaf, of bent grasses caused by **H. erythrospilum,** n. sp., compared with spot due to **H. triseptatum,** 349–350; leaf, see leaf spot
Spotted wilt, on truck and ornamental plants in California, 17; transmission, 17
SPRAGUE, RODERICK, Ascochyta boltshauseri on beans in Oregon, 416-420; Wojnowicia graminis as a very weak, secondary parasite of winter cereal crops, 405–415
Spray, oil-soluble copper, use of, as fungicide, 368–371
Spraying, and dusting for control of sugar beet leaf spot in Southern Minnesota, 234–243; for apple scab, in relation to leaf growth rate, 373–375
Sprays, fungicidal, containing copper, penetration of plant tissues by, 368–370; peach, copper fungicides, etc., 972
Squash, sclerotinia rot in Montana, 39–40; spread of Southern celery-mosaic virus on, in Florida, 289–308
Stability of antisera, 697
Stagonospora, crini, synonym of S. curtisii, 266; curtisii, inoculation of, on Amaryllidaceae, 262–268; curtisii, on Crinum powelli, 263; curtisii, Phyllosticta hymenocallidis, a synonym of, 266; curtisii, S. crini, a synonym of, 266
Stagonosporopsis, boltshauseri, synonym of Ascochyta boltshauseri, 419; hortensis, synonym of A. boltshauseri, 419; invalidity of genus, 419
Stagonopsis phaseoli, synonym of Ascochyta boltshauseri, 419
STAKMAN, E. C., see CHRISTENSEN, J. J.; see LEVINE, M. N.; LEE HINES, ROBERT C. CASSELL, and M. N. LEVINE, Population trends of physiologic forms of Puccinia graminis tritici, 1930 to 1934, 34; LEON J. TYLER, G. E. HAFSTAD and E. G. SHARVELLE, Experiments on physiologic specialization and nature of variation in Ustilago zeae, 34
Staling products, effect of, on growth of Coccomyces hiemalis, 141, 142
STANLEY, A. R., Studies on the soft-rot and colon-typhoid-dysentery groups of bacteria. 2, Physiology, 34–35
STANLEY, W. M., Chemical studies on the virus of tobacco mosaic, 475–492; Chemical studies on the virus of tobacco mosaic. IV. Some effects of different chemical agents on infectivity, 899–921; Chemical studies on the virus of tobacco mosaic. V. Determination of optimum hydrogen-ion concentration for purification by precipitation with lead acetate, 922–930
Starvation of tobacco due to lack of sulphur, 430–432
STEINMETZ, F. H., and M. T. HILBORN, A method of determining the rate of decay in wood, 35
Stellaria media, observations on Ligniera junci, a root parasite of, 893
Stem, blight, of Lespedeza sericea, 976; -infesting nematode, new hosts of, 1003–1010; rot, sweet potato, control of, 976; rust, decrease in severity, factors responsible, 25; rust, decrease in severity, indicated by uniform rust nurseries, 25; rust,

prevalence of physiologic forms, 34; rust, wheat, carotenoid pigments of urediospores, 30; rust, wheat, color variants, of, 30

Sternbergia lutea, infection with Staganospora curtisii, 264

STEVENS, NEIL E., see MACKIE, W. W.; and WOOD, JESSIE I., Losses from corn ear rots in the United States, 281–283

STEVENSON, J. A., see CLAYTON, E. E.

Stewart's disease of corn, bacteriophage in relation to, 371–372; host range, 992–1002

Stilbum, cinnabarinum, ascigerous stage of, 215–217; cinnabarinum, description of, 213–217; cinnabarinum, pathogenicity of, 217, 218; cinnabarinum, relation to host, 219; disease of fig, 208–222

STONE, G. M., Control of sweet-potato stem rot in West Tennessee, 976; see SHERBAKOFF, C. D.

Strawberries, Sclerotium rolfsii on, and effect of certain chemicals on sclerotia, 966

Strawberry, anthracnose and wilt caused by Colletotrichum fragariae, 973; black stele root rot, 5; diseases, review of papers on, see Darrow, G. M., 523–525; Phytophthora, 5; root rot of, caused by species of Diplodia, 974–975

Streak, British Queen, serological evidence in classification of virus, 687–700; maize, retention of virus by aphids, 175; possible source of infection in black raspberries, 959–961; rose, transmission of virus, 7–8; virus disease of roses, 7–8

STROMAN, G. N., J. J. TAUBENHAUS, and WALTER N. EZEKIEL, Some effects of phymatotrichum root rot on the microscopic characters of cotton fibers, 126–130

STUART, W. W., and A. G. NEWHALL, Further evidence of the seed-borne nature of Peronospora destructor, 35

Studies, cultural, on the virus of tobacco mosaic, 1035–1037; on frenching of tobacco, 1067–1084; on seven differentiating characteristics of two physiologic forms of Tilletia tritici, 1091–1098; on the soft-rot and colon-typhoid-dysentery-groups of bacteria. 1. Pathogenicity, 19; on the soft-rot and colon-typhoid-dysentery groups of bacteria. 2. Physiology, 34–35

Study of coalescing haploid pustules in Puccinia helianthi, 1085–1090

Sudan grass, infection with Phytomonas stewarti, 996–997; inoculation with P. stewarti, 21

Sugar, beet bacterial-pocket disease, physiologic studies of bacteria, 31; beet, leaf spot control in Southern Minnesota, 234–243; beets, damping off, influence of preceding crops on, 13; beets, factors affecting experimental error in greenhouse pot tests, 1019–1025; beets, root rot of, by Sclerotium rolfsii, 957–959; cane, inoculations with Phytomonas stewarti negative, 997; cane, mosaic, severe type of, 36–37; cane, root diseases, relation of insects to, 12

Sugars, effect on variability of Coccomyces hiemalis, 148

Sulphur, aeroplane dusting with, to combat stem rust of wheat, 892–893; as fungicide, studies on, 35–36; soil treatment with, 16; starvation, in tobacco, 430–432

SUMNER, C. B., see ARMSTRONG, G. M.

Susceptibility, influence of phosphorus and potassium supply on, to yellow tobacco mosaic infection, 493–502; of host to nitrogen supply, effect on virus infection, 178–191; of tissue to mosaic virus infection, 188, 189; to yellow tobacco mosaic, influence of nutrition on, 33–34

Sweet, corn, inoculation with Phytomonas vascularum, 21; corn, resistant to bacterial wilt, 21–22; corn, spread of Southern celery-mosaic virus on, 289–308; pea, fasciation, due to bacteria, 36; pea fasciation, seed transmission of organism, 36; pea, infected with Anguillulina dipsaci, 1006; potato, control of stem rot of, 976

Symbionts of Pseudococcus brevipes in relation to a phytotoxis secretion of the insect, 10

Symptoms, of narcissus mosaic developed within the plant, 896–897; of plant viruses, 330–343; of rose mosaic, 8

Tagetes, erecta, susceptible to celery virus 1, 386, 398; patula, susceptible to virus 1, 386, 398

Tannic acid, effect on infectivity of tobacco-mosaic virus, 931–937

Tape, antiseptic adhesive, in control of hairy root and crown gall, 192–207

Taphrina, deformans, copulation of conidia, 56–57; deformans, cytology, 56; deformans, development of mycelium in the host, 54–55; deformans, hibernation of conidia, 50–51; deformans, inoculation experiments with monosporous cultures, 59–61; deformans, life history of, 41–66; deformans, nucleation of cells formed in culture, 61–63; deformans, occurrence of late infection, 49–50; deformans, origin of saprophytic phase, 63–64; deformans, penetration into host, 52–53; deformans, perennial mycelium, 51–52; deformans, spraying against, 48; epiphylla, copulation of conidia, 58; klebahni, copulation of conidia, 58

TAPKE, V. F., An effective and easily applied method of inoculating seed barley with covered smut, 1038–1039

Tarsonemus latus, mite, on potatoes, beans, and apples, 654

TAUBENHAUS, J. J., On a black crown rot of greenhouse snapdragons caused by Myrothecium roridum, 969–970; Seeds of watermelons and okra as possible carriers of fusarium wilt, 969; see EZEKIEL, WALTER N.; see STROMAN, G. N.; and PHARES DECKER, Laboratory and field studies on sulphur as a fungicide, 35–36; and W. N. EZEKIEL, Further studies on the longevity of sclerotia of Phymatotrichum omnivorum, 36; and WALTER N. EZEKIEL, The quality of lint and seed from cotton plants with Phymatotrichum root rot, 104–113

Taxonomy of the Fusaria that cause celery wilt (yellows , 29

Technique for isolation of Ceratostomella ulmi from bark beetles, 37–38

Temperature, effect of on development of **Helminthosporium erythrospilum**, n. sp., cause of leaf spot of bent grasses, 347–349; effect of, on diameter of colonies of Coccomyces hiemalis, 143–144; effect of, on infection of Hope wheat by Puccinia graminis tritici, 1041–1066; for demonstrating Fusarium in barley seed, by germination, 313–315; for demonstrating Helminthosporium in barley seed, by germination, 313–315; high, effect of, on infection by Puccinia graminis tritici, 1058–1059; relation to growth of Fusarium causing celery yellows, 583–585; soil, effect of, on development of celery yellows, 585–590; tank, Chicago soil-nutrient, suitability for phytopathologic studies, 26

Teosinte infected by Aplanobacter stewarti, 32

Tetrachlorethane, effect on phymatotrichum root rot and host plants, 16

Tetragonia expansa, susceptible to celery virus 1, 381, 385, 395

Thalictrum spp., inoculation experiments, with Puccinia rubigo-vera and P. tomipara, 663–679

Thelocactus nidulans, disease of, 9

Thermal, death point, of plant viruses, 330–343; inactivation of virus, and precipitin reactivity, 703

Thielaviopsis basicola, albino culture of, 1014–1016; cultural variations of, 1011–1018; hyphal characters, 1016; no perfect stage, 1016; pathogenicity tests, 1016; single-spore cultures, 1011–1014; spore variations, 1016

Thistle, Canada, sclerotinia wilt of, 963–964

THOMAS, R. C., A bacteriophage in relation to Stewart's disease of corn, 371–372

THORNBERRY, H. H., Effect of phosphate buffers on infectivity of tobacco-mosaic virus, 618–627; Effect of tannic acid on the infectivity of tobacco-mosaic virus, 931–937; Particle diameter of certain plant viruses and Phytomonas pruni bacteriophage, 938–946; Particle size of three strains of tobacco-mosaic virus, 36; Quantitative studies on the filtration of tobacco-mosaic virus, 601–617; see REED, H. S.

THORNTON, J. K., Blackberries: Possible source of streak infection in black raspberries, 959–961

Three forms of the fusarium wilt of celery, 29

Thrips tabaci, transmission of spotted wilt virus by, 17

Thuja spp., crown gall on, 894

TILFORD, P. E., Fasciation of sweet peas: A bacterial disease, 36; The use of the quinhydrone and antimony electrodes for determining the pH of solid culture media, 362–367

Tilletia, levis, artificial infection of rye (var. Petkus) 503–515; secalis, occurrence in Argentina, 513; tritici, artificial infection of rye, 503–515; tritici, new variety of, in Montana, 40; tritici, physiologic forms of, differentiating characteristics of, 1091–1098

TIMS, EUGENE C., Severe type of mosaic on a sugar-cane variety, 36–37; A Stilbum disease of figs in Louisiana, 208–222

TISDALE, W. B., A bark disease of Tahiti lime trees caused by Phomopsis citri and Diplodia natalensis, 966

To visualize a distinction between viruses and organisms, 892

Tobacco, Ambalema, resistance to different viruses, 37; bed soil disinfection, 972; black-shank, soil temperature as factor in development of, 975; calomel, injurious to root system, 6; downy mildew, caused by Peronospora tabacina, 516–521; downy mildew, nomenclature of fungus, 516–521; downy mildew of, control of, 19, 971; effect of certain environmental factors on germination of Florida cigar-wrapper seeds, 975; effect of nitrogen on susceptibility to yellow tobacco virus, 184–188; example of spread of veinbanding to, from potatoes, 650–652; frenching, deficiency studies on, 1074–1081; frenching, effect of, on development of and recovery from, of aluminum, 1075–1078; frenching effect of, on development of and recovery from, of copper, 1075; frenching, effect of soil aeration on, 1073; frenching, effect of soil reaction on, 1073–1074; frenching, effect of soil texture on, 1072–1073; frenching, not due directly to mineral deficiency, 1074–1081; frenching, not due to parasite, 1071–1072; frenching, parasitic studies on, 1071–1072; frenching, physiologic studies on, 1072–1082; frenching, probably a toxicity disease, 1082; frenching, studies on, 1067–1084; frenching, symptoms of, 1068–1070; frenching, toxicity studies on, 1081–1082; incidence of downy mildew as related to air-borne sporangia, 636; incidence of downy mildew as related to seed-bed practices, 636–638; incidence of downy mildew as related to site of seed bed, 630–636; initiation of downy mildew in North Carolina in 1934, 628–639; leaf diseases, epidemiology, effect of, driving rains, 11; leaf diseases, epidemiology, effect of fertilizers, 11; leaf diseases, epidemiology, effect of topping, 11; mosaic, causing mosaic burn, 967; mosaic, chemical studies of virus, 475–492; mosaic, chemical studies on the virus of, 899–921; mosaic, cultural studies on the virus of, 1035–1037; mosaic, determination of optimum hydrogen-ion concentrations for purification by precipitation with lead acetate, 922–930; mosaic, effect of phosphate buffers on infectivity of virus, 618–627; mosaic, effect of tannic acid on infectivity of, 931–937; mosaic, longevity of virus, factors affecting, 15; mosaic, quantitative studies on filtration of virus, 601–617; mosaic, strains, serological evidence in classification of viruses, 687–700; -mosaic virus, effect of proteoclastic enzymes on, 33; -mosaic virus, particle size of three strains of, 36; -mosaic viruses, relation of necrotic lesions to localization of, in Nicotiana, 968; ring spot, inactivation of virus at different hydrogen-ion concentrations, 487–489; ring spot, longevity of virus, factors affecting, 15; ring spot, recovery and immunity from, 37; soil-temperature studies on cigar-wrapper, 975; sulphur starvation in, 430–432; symptoms of crucifer virus on, 640–642; viruses,

thermal inactivation of, 15; wildfire, control of, in Pennsylvania, 6; wildfire, on Physalis virginiana, 6; yellow mosaic, influence of nutrition on susceptibility, 33–34; yellow mosaic, susceptibility to, influenced by phosphorus and potassium, 493–502

Tomato, "combination-streak," virus complex, chemical splitting of, 33; "combination-streak," effect of certain chemicals on virus of, 864–874; graft *versus* insect transmissions of curly top (tomato yellows) in, 844–853; late blight, injurious outbreaks, in Massachusetts, 7; late blight, seed-borne, 7; pathologico-anatomical changes in, incident to woodiness of fruit, 539–558; seedlings, control of diseases of, 968

TOMPKINS, C. M., C. M. TUCKER and A. E. CLARKE, Root rot of aster caused by Phytophthora cryptogea, 895; C. M. TUCKER, and M. W. GARDNER, A Phytophthora root rot of cauliflower, 893–894; see GARDNER, M. W.

Top-necrosis virus, found in apparently healthy potatoes, 1115–1116

Toxic effect on pineapple of feeding by Pseudococcus brevipes, 10

Toxicity, of compounds of ammonia to Sclerotium rolfsii, 895–896; of copper-lime-arsenite preparations, 23–24; of cuprous ion as compared with cupric, 21; of eradicant fungicides, 23–24; of plant juice, reduced by dialysis, 689; studies, on tobacco frenching, 1081–1082

Tradescantia sp., susceptible to celery virus 1, 392, 394

Tragopogon porrifolius, infected with Anguillulina dipsaci, 1003–1005

Trametes lactinea, secondary spore formation in, 426

Transfer of Diaporthe umbrina to the genus Cryptosporella, 886–889

Transmission, of cabbage mosaic, 6; of Ceratostomella ulmi through root grafts, 1039–1040; of plant viruses, modes of, 330–343

Treatment, cotton seed, results from ten years' work, 972; of barley seed for control of seedling blight, 318–326; of corn seed, with bacteriophage, for control of Stewart's disease, 371–372

Treatments, cottonseed, in South Carolina, 970

Trees, fruit, little leaf or rosette in California, 275–278; fruit little leaf or rosette, see CHANDLER, W. H., 522

Trichoderma, antagonistic action of, on Rhizoctonia and other soil fungi, 244–252; lignorum, inhibiting effect on Phymatotrichum omnivorum, 573–574; toxicity of filtrate affected by heat, 247–249; toxicity of filtrate to Pythium de baryanum, 251; toxicity of filtrate to Rhizoctonia, 250, 251

Trifolium incarnatum, infection with pea mosaic, 162, 163

Triticum, aestivum, susceptible to celery virus 1, 392, 393, 394; dicoccum, Wojnowicia graminis on, 408; monococcum, W. graminis on, 408; spelta, W. graminis on, 408

Tropaeolum majus, susceptible to celery virus 1, 386, 395

Trunk canker, of apple, caused by Phytophthora cactorum, 5; of apple, control, 5; of apple, destructive disease, 5; of apple, symptoms, 5; of apple, varietal susceptibility to, 5

Trypsin, effect of, on tobacco-mosaic virus, 33

TU, CHIH, and LI, H. W., Breeding millet resistant to smut in North China, 648–649

Tuberculina, maxima, as control means for Cronartium species, 259, 260; maxima, growth on artificial media, 256; maxima, inoculations attempted, 256; maxima, observations on, a parasite of Cronartium ribicola, 253–261; maxima, on aecia of C. cerebrum, 1117–1118; maxima, spore description, 255; persica, synonym of T. maxima, 253, 255

TUCKER, C. M., see TOMPKINS, C. M.

Turnip, roots of, rotted by Macrosporium and Colletotrichum, 269–274; virus disease of, 640

TYLER, LEON J., see STAKMAN, E. C.; and CLYDE P. SHUMWAY, Hybridization be-

tween Sphacelotheca sorghi and Sorosporium reilianum, 375–376

Ulmus, parvifolia, Chalaropsis thielairoides on, 652–654; pumila, C. thielairoides on, 652–654

Ultra-violet light, effect on conidial production of Coccomyces hiemalis, 144, 145

UNDERWOOD, J. K., Lespedeza sericea stem blight, 976

Unidentified species of Sphaeropsis on maize in California, 893

Uredinopsis mirabilis, host of Tuberculina maxima, 255

Uromyces appendiculatus, effect of light on infection by, 715–725

Use, of oil-soluble copper as a fungicide, 368–370; of the quinhydrone and antimony electrodes for determining the pH of solid culture media, 362–367

Ustilago, agrostis-palustris, 816; clintoniana, on Dactylis glomerata, 815, 816; crameri, on Chaetochloa italica in North China, 648–649; hordei, 28; hordei, albino strain, monosporidial lines, compatible with normal Ustilago hordei and U. medians, 27–28; hordei, albino strain of, 27–28; hordei, albino strain, sexual groups, 28; hordei, hybridization between U. hordei and U. medians, 5; hordei, method of inoculating seed barley with, 1038, 1039; hordei, pathogenicity, 5; medians, 28; medians, distribution of, 28; medians, hybridization between U. medians and U. hordei, 5; medians, pathogenicity, 5; nuda, color groups of, 28; phlei-pratensis, 816; poae-annuae, 816; poae-pratensis, 816; striaeformis, forma agrostidis, on Agrostis palustris, 816; striaeformis, forma phlei, on Phleum pratense, 816; striaeformis, forma poae-annuae, on Poa annua, 816; striaeformis, forma poae-pratensis, on Poa pratensis, 816; striaeformis, on some common grasses, 810–817; zeae, effect of infections on yield of corn, 223–233; zeae, nature of variation in, 34; zeae, physiologic specialization in, 34

VALLEAU, W. D., A probable case of sulphur starvation in tobacco, 430–432; Do necrotic lesions result in localization of tobacco-mosaic viruses in Nicotiana? 968; Do tobacco plants recover and develop an immunity from ring spot? 37; The resistance of Ambalema tobacco to different viruses, 37; Seed transmission of Helminthosporium of corn, 1109–1112; see JOHNSON, E. M., and E. M. JOHNSON, Only certain strains of tobacco mosaic cause mosaic burn, 967

VAN HALTERN, FRANK, Control of diseases of tomato seedlings, 968

Variability in monosporic cultures of Coccomyces hiemalis, 131–159

Variation, discontinuous, in Coccomyces hiemalis, 155–156; in Ustilago zeae, 34

Variations, cultural, of Thielaviopsis basicola, 1011–1018

Varietal susceptibility of apple to trunk canker, 5

Vascular disease in poplar and willow, 18–19

Vector, curly-top, effect of carbon dioxide on the feeding of, 891

Vectors, insect, of plant viruses, 330–343

Vegetable crops, diseased by Southern celery-mosaic virus, 380–385

Vegetables, dissemination of Southern celery-mosaic virus on, in Florida, 289–308

Veinbanding, example of spread from potatoes to tobacco, 650–652; of potato, serological evidence in classification of virus, 687–700

Venturia, inaequalis, control by eradicant fungicide, 23; inaequalis, leaf growth rate in relation to control of, 373–375; pyrina, control by eradicant fungicide, 23

Verbena, host for spotted wilt in California, 17

Vermicularia, polytricha, synonym of **Colletotrichum truncatum**, n. comb., 121, 122; truncata, synonym of **Colletotrichum truncatum**, n. comb., 121, 122; sp., 886

VERRALL, A. F., and T. W. GRAHAM, The transmission of Ceratostomella ulmi through root grafts, 1039–1040
Verticilliosis, experiment to determine susceptibility of flax to, 892
Verticillium, albo-atrum, susceptibility of flax to, 892; strains of, 972; wilt disease of cotton, 972
Vicia faba, mosaic, 160–177; susceptible to celery virus 1, 392, 395; vehicle of immunization antigen, 689
Vigna sinensis, description of cucumber-mosaic virus on, 950; root-knot nematode on, 72–74; var. Black Eye (Cowpea), experimental material, infectivity of tobacco-ring-spot and cucumber-mosaic viruses, 487–489
Vinca rosea, susceptible to celery virus 1, 386, 396
Virus, acquired immunity in zinnia from cucumber-mosaic, 776–789; as actual antigenic substance, 709; aucuba-mosaic, symptoms in zinnia, 779; aucuba mosaics, particle diameter of, 938–946; celery 1, host plants, 381, 386, 389, 392, 394–398; chemical combination with silver, 707; "combination-streak," tomato, components of, 864; "combination streak" of tomatoes, effect of certain chemicals on, 864–874; Commelina nudiflora, symptoms compared with yellow-spot, on pineapple, 10; common bean mosaic, inheritance of resistance to, 875–883; cucumber-mosaic, similarity of Southern celery-mosaic to, 947; cucumber-mosaic, symptoms in zinnia, 777; curly-top, experiments on the multiplication of, in the beet leaf hopper, 891; curly-top, not transmitted by seed, 894; disease, streak, on roses, 7–8; diseases of plants, D. ATANASOFF, review by JAMES JOHNSON, 444; effect of dilution on filterability of tobacco-mosaic, 604; effect of hydrogen-ion concentration on filterability of tobacco-mosaic, 605; effect of nitrogen supply on host susceptibility of, 178–191; effect of phosphate buffers on infectivity of tobacco-mosaic, 618–627; fractional filtration of tobacco-mosaic, 606–609; inactivation at different hydrogen-ion concentrations, cucumber-mosaic, 487–489; inactivation at different hydrogen-ion concentrations, tobacco ring-spot, 487–489; longevity of tobacco-mosaic and ring-spot, factors affecting, 15; mosaic of peas, retention by aphid, 170–174; of crucifers and other hosts, 640–644; of pea mosaic, incubation of, in aphids, 31; of tobacco mosaic, determination of optimum hydrogen-ion concentrations for purification by precipitation with lead acetate, 922–930; of tobacco mosaic, effects of different chemical agents on infectivity, 899–921; peach rosette, destroyed by heat treatment of trees, 24; peach yellows, destroyed by heat treatment of trees, 24; potato latent mosaic, component of tomato "combination-streak," 864; potato ring-spot, particle diameter of, 938–946; precipitin reaction of, not affected by host, 709; properties of crucifer mosaic, 642–643; severe-etch, symptoms in zinnia, 780; size of particle of tobacco-mosaic, 613; Southern celery-mosaic, a strain of cucumber-mosaic virus, 953; Southern celery-mosaic, causal factors in increase of, 301–304; Southern celery-mosaic, classification of, 947–954; Southern celery-mosaic, dissemination of, on vegetable crops in Florida, 289–308; Southern celery-mosaic, factors influencing field dissemination of, 304–306; Southern celery-mosaic, host range of, 377–404; Southern celery-mosaic, insect vector of, in Florida, 290–308; Southern celery-mosaic, natural history of, 306–307; Southern celery-mosaic, spread of, in seed beds, 302; Southern celery-mosaic, wild hosts of, in Florida, 290–308; spotted wilt of tomato, 898; spotted wilt on truck and ornamental plants, 17; spotted wilt, transmission, 17; spotted wilt, transmission, by Frankliniella sp., 17; spotted wilt, transmission by Thrips tabaci, 17; streak, on roses, symptoms, 8; streak, on roses, transmission of, 8; susceptibility to, influence of phosphorus and potassium supply on, 493–502; tobacco-

mosaic, chemical studies on, 475–492; tobacco-mosaic, cultural studies of, 1035–1037; tobacco-mosaic, effect of proteoclastic enzymes on, 33; tobacco-mosaic, effect of tannic acid in infectivity of, 931–937; tobacco-mosaic, green, component of tomato "combination-streak," 864; tobacco-mosaic, longevity of, factors affecting, 15; tobacco-mosaic, necrotic type, symptoms in zinnia, 779; tobacco-mosaic, particle size of three strains of, 36; tobacco-mosaic, quantitative studies on filtration of, 601–617; tobacco-mosaic, rates of inactivation at different hydrogen-ion concentrations, 475–492; tobacco-mosaic, symptoms in zinnia, 778; tobacco mosaics, particle diameter of, 938–946; tobacco-ring-spot, longevity of, factors affecting, 15; tobacco-ring-spot, particle diameter of, 938–946; tobacco-ring-spot, symptoms in zinnia, 780; tobacco-ring-spot, yellow, symptoms in zinnia, 780; tomato "combination-streak," complex chemical splitting of, 33; top-necrosis, from apparently healthy potatoes, 1115–1116; top-necrosis, from apparently healthy potatoes, same as virus B, 1116; ultrafiltration of tobacco-mosaic, 612–614; undetermined, of tomato, 898

Viruses, and organisms, distinction between, 892; antigenicity of plant, 702–714; cause of Oregon tip blight of tomato, 897–898; classification of serological evidence in, 686–701; classification of value of specific immune reaction in, 776; complement-fixation tests with, 691; effect of progressive inactivation on precipitin reactivity, 703–707; Mez reaction, 691–692; on pineapple, mechanical transmission, 10; plant, bases for description and identification of, 329–343; plant, chief groups of, 332; plant, descriptive key for, 328–343; plant, incompletely described, list of, 335–336; plant, insect vectors of, 329–343; plant, longevity *in vitro* of, 329–343; plant, modes of transmission of, 329–343; plant, natural and differential hosts of, 329–343; plant, particle diameter of, and Phytomonas pruni bacteriophage, 938–946; plant, symptoms of, 329–343; plant, thermal death point of, 33–343; precipitin testing, 689–691; preparation of immune sera, 688–689; resistance of Ambalema variety to, 37; serological evidence of relationships, 10; serological reaction of, specificity of, 697; tobacco-mosaic, relation of necrotic lesions to localization of, in Nicotiana, 968; tobacco, thermal inactivation of, 15; tulip-breaking, properties and interpretations of, 898

WALTER, J. M., Technique advantageous for the isolation of Ceratostomella ulmi from bark beetles, 37–38

WARE, J. O., see YOUNG, V. H.

Washingtonia filifera, Penicillium vermoeseni on, in California, 896

Watermelon, susceptibility to Phymatotrichum omnivorum, 559–577

Watermelons, seeds of, as possible carriers of fusarium wilt, 969

Weather, effect of, on development of **Helminthosporium erythrospilum**, n. sp., cause of leaf spot of bent grasses, 347–349; influence of phytophthora disease of peach seedlings, 802

WEBER, G. F., An aerial Rhizoctonia on beans, 38; The direct effect of Bordeaux mixture on early cucumber production, 974

Weeds, hosts of celery virus, 388; Southern celery mosaic on, in Florida, 289–308

WEHMEYER, LEWIS E., see JENKINS, ANNA E.

WEISS, FREEMAN, A fungus spot of azalea flowers, 38; and THELMA B. POST, Bottom rot of calla rhizones, 38–39

WELLMAN, F. L., Dissemination of Southern celery-mosaic virus on vegetable crops in Florida, 289–308; The host range of the Southern celery-mosaic virus, 377–404

Wheat, aeroplane dusting with sulphur to combat stem rust of, 892–893; Hope, effect of extract from sap on germination of urediospores of Puccinia graminis tritici, 1049–1050; Hope, effect of other organisms on development of stem-

rust infection in, 1053–1054; Hope, infection by P. graminis tritici, effect of environment on, 1054–1055; Hope, infection by P. graminis tritici, effect of high temperature on, 1058–1059; Hope, infection by P. graminis tritici, effect of light intensity and temperature on, 1041–1066; Hope, infection by P. graminis tritici, effect of reduced light intensity on, 1055–1058; Hope, inoculation with stem rust, by hypodermic syringe, 1050–1053; Hope, occurrence of stem rust on, in 1933, 1045–1047; Hope, resistance of, to stem rust, 1045; Hope, resistance of, to stem rust of older plants, 1047–1049, 1053; Hope, susceptibility of, to stem rust, of young plants, 1047; infection with P. graminis, effect of light on, 715–725; infection with P. triticina, effect of light on, 715–725; physiologic forms of Tilletia tritici on, 1091–1098; seedlings, susceptibility of, to P. graminis, 1044; stem rust, and color variants of, study of carotenoid pigments of urediospores of, 30; stem rust, color variants of, 30; T. tritici on, new variety of, in Montana, 40; winter, Wojnowicia graminis on, 407

WHIPPLE, O. C., see GARDNER, M. W.

White-bud disease of raspberry, caused by Glomerella rubicola, 93; cultural character of organism, 93–95; economic importance, 93; host relationships, 92–93; relation to bitter rot of apples, 93–100; symptoms, 91–92

WILCOX, HAZEL W., and GEORGE K. K. LINK, Serically active (haptenic) carbohydrates of genotypes of Neurospora tetrasperma and N. sitophila, 39

Wildfire, control, 6; on Physalis virginiana, 6; on tobacco, in Pennsylvania, 6

Willow, vascular disease, 18–19

WILSON, A. R., The influence of Phytomonas tumefaciens and P. rhizogenes on the actual acidity of certain liquids and agar substrata, 854–863

Wilt, cotton, reaction of fungus to toxic dyes, 970; disease, corn (Stewart's) host range, 992–1002; fusarium, control of, in cotton, 969; fusarium, of celery, forms of, 29; fusarium, of cotton, 975; fusarium, seeds of watermelons and okra as possible carriers of, 969; of celery, taxonomy of Fusaria causing, 29; of China aster due to Fusarium, 733–747; of corn, Stewart's disease, control of, by treatment of seed with bacteriophage, 371–372; of strawberry, caused by Colletotrichum fragariae, 973–974; -resistant cottons adapted to the Gulf Coastal Plains, 974; sclerotinia, of Canada thistle, 963–964; verticillium disease of cotton, 972

WINGARD, S. A., Host-parasite relationship in bean rust, 39

Wojnowicia, graminis, a weak, secondary parasite of winter cereal crops, 405–415; graminis, associated with Cercosporella herpotrichoides, 406, 408; graminis, host range, 407–408; graminis, inoculation studies with, on cereals and grasses, 408–410; graminis, taxonomy of, 410–413; hirta, comparison with W. graminis, 412; tenella, comparison with W. graminis, 412

WOLF, FREDERICK A., see DIXON, LAWRENCE F.

Wood, decay, determination of rate of, 35

WOOD, JESSIE I., see STEVENS, NEIL

WRIGHT, ERNEST, see LAMB, HOWARD

Xanthium americanum, Rhizoctonia sp. on, in Florida, 38

Xylaria mali, isolated from apple roots, 976; on apple, relation of host vigor, to infection by, 12–13

Xylol, effect on phymatotrichum root rot and host plants, 16

Yeast on pine attacked by Southern pine beetle, 7

Yellow foxtail, infection of, with Phytomonas stewarti, 997

Yellows, and rosette of peach, heat treatment cure, 24; Fusarium, of celery, 578–600; or fusarium wilt of celery, 29; psyllid, observations on pathological histology and phyto-chemistry of, 895

YOUNG, H. C. and J. R. BECKENBACH, Insoluble copper compounds as substitutes for Bordeaux, 40

YOUNG, P. A., A new variety of Tilletia tritici in Montana, 40; Sclerotinia rot of pumpkin and squash, 39–40

YOUNG, V. H., J. O. WARE, and O. A. POPE, Control of potash hunger and fusarium wilt in cotton, 969

ZALESKI, KAROL, see HART, HELEN

Zantedeschia aethiopica, bottom rot of rhizomes, 39

Zea mays, infection with Phytomonas vascularum, 997–998; infection with Puccinia sorghi, effect of light on, 715–725; pathologic significance of seed-coat injury, 24; resistant to bacterial wilt, 21–22; see corn; Stewart's wilt disease of, host range, 992–1002; susceptible to celery virus 1, 381, 383, 394; transmission of mosaic viruses to, 952

Zebrina pendula susceptible to celery virus 1, 386, 387, 394

Zephyranthes candida, infection with Stagonospora curtisii, 264

Zinc sulphate for peach spray, 22–23

Zinnia, elegans, acquired immunity from cucumber mosaic, 776–789; elegans, effect of virus test on, 947; elegans, susceptible to celery virus 1, 386, 387, 398; viruses, symptoms produced by, 777–780

Zone phenomenon in complement-fixation tests, 691

Zoosporangia production in Phytophthora, stimulated by nonsterile soil leachate, 432–435